

ISOGENY CLASSES OF ABELIAN VARIETIES OVER FUNCTION FIELDS

YURI G. ZARHIN

ABSTRACT. We study finiteness problems for isogeny classes of abelian varieties over an algebraic function field K in one variable over the field of complex numbers. In particular, we construct explicitly a non-isotrivial absolutely simple abelian fourfold X over a certain K such that the isogeny class of $X \times X$ contains infinitely many mutually non-isomorphic principally polarized abelian varieties. (Such examples do not exist when the ground field is finitely generated over its prime subfield.)

2000 Mathematics Subject Classification 14K02 (primary), 14D07 (secondary)

1. INTRODUCTION

Let K be a field, $\bar{K}$ its separable closure, $\text{Gal}(K) = \text{Gal}(\bar{K}/K)$ the (absolute) Galois group of K . Let X be an abelian variety over K , $\text{End}_K(X)$ its ring of K -endomorphisms and $\text{End}_K^0(X) := \text{End}_K(X) \otimes \mathbf{Q}$.

Let us consider the set $\text{Isog}(X, K)$ of K -isomorphism classes of abelian varieties Y over K such that there exists a K -isogeny $Y \rightarrow X$. If ℓ is a prime different from $\text{char}(K)$ then let us consider the set $\text{Isog}(X, K, \ell)$ of K -isomorphism classes of abelian varieties Y over K such that there exists an ℓ -isogeny $Y \rightarrow X$ that is defined over K . (Recall that ℓ -isogeny is an isogeny, whose degree is a power of ℓ .) If m is a positive integer then we write $\text{Isog}_m(X, K, \ell)$ for the subset of $\text{Isog}(X, K, \ell)$ that consists of all (isomorphism) classes of Y with a K -polarization of degree m .

When K is finitely generated over its prime subfield, Tate [32] conjectured the finiteness of $\text{Isog}_m(X, K, \ell)$. This conjecture played a crucial role in the proof of Tate's conjecture on homomorphisms and the semisimplicity of the Galois representations in the Tate modules of abelian varieties [32, 28, 24, 33, 34, 35, 10, 11, 20, 39].

If K is finitely generated over the field $\mathbf{Q}$ of rational numbers then it was proven by Faltings [11] that $\text{Isog}(X, K)$ is finite. (See also [10, 38].) When K is finite, the finiteness of $\text{Isog}(X, K)$ is proven in [36]. (See [24, 33, 35, 36, 20] for a discussion of the case when K is infinite but finitely generated over a finite field.)

The aim of this note is to discuss the situation when K is an algebraic function field in one variable over $\mathbf{C}$. (In other words, K is finitely generated and has transcendence degree 1 over $\mathbf{C}$.) In order to state our results, notice that one may view K as the field of rational functions on a suitable irreducible smooth algebraic complex curve S such that X becomes the generic fiber of an abelian scheme $f : \mathcal{X} \rightarrow S$. Let $s \in S$ be a complex point of S , let $\mathcal{X}_s$ be the fiber of f at s and $H_1(\mathcal{X}_s, \mathbf{Q})$ its first rational homology group. (Recall that $\mathcal{X}_s$ is a complex abelian variety, whose dimension coincides with $\dim(X)$.) There is the natural

global monodromy representation

$$\pi_1(S, s) \rightarrow \text{Aut}(H_1(\mathcal{X}_s, \mathbf{Q}))$$

of the fundamental group $\pi_1(S, s)$ of S . Deligne [6] proved that this representation is completely reducible and therefore its centralizer $D_f := \text{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Q}))$ is a (finite-dimensional) semisimple $\mathbf{Q}$ -algebra. On the other hand, every $u \in \text{End}_K(X)$ extends to an endomorphism of $\mathcal{X}$ and therefore induces a certain endomorphism u_s of $\mathcal{X}_s$. This gives rise to the embeddings

$$\text{End}_K(X) \hookrightarrow \text{End}(\mathcal{X}_s) \hookrightarrow \text{End}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q})),$$

whose composition extends by $\mathbf{Q}$ -linearity to the embedding

$$\text{End}_K^0(X) \hookrightarrow \text{End}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q})),$$

whose image lies in D_f . Further, we identify $\text{End}_K^0(X)$ with its image in D_f . Our main result (Theorem 4.1) may be restated as follows.

The set $\text{Isog}(X, K)$ is infinite if and only if $D_f \neq \text{End}_K^0(X)$.

If $D_f \neq \text{End}_K^0(X)$ then the set $\text{Isog}(X, K, \ell)$ is infinite for all but finitely many primes ℓ (see Theorem 4.3). In addition, if X is principally polarized over K then the set $\text{Isog}_1(X^2, K, \ell)$ is infinite for all but finitely many primes ℓ congruent to 1 modulo 4 (see Corollary 4.6).

In order to describe other results of this paper (Sect. 10), let us further assume that every homomorphism over $\overline{\mathbf{C}(S)}$ between X and “constant” abelian varieties (defined over $\mathbf{C}$) is zero. If $\dim(X) \leq 3$ then Deligne [6] proved that $D_f = \text{End}_K^0(X)$: this implies that in this case $\text{Isog}(X, K)$ is finite. On the other hand, Faltings [9] constructed a four-dimensional X with $D_f \neq \text{End}_K^0(X)$; in his example(s) X is the generic fiber of an universal family of abelian fourfolds with level $n \geq 3$ structure over a Shimura curve.

We prove that if $\dim(X) = 4$, all endomorphisms of X are defined over K and $\text{Isog}(X, K)$ is infinite then $\text{End}_K^0(X)$ is a CM-field of degree 4 (see Theorem 10.1). Almost conversely, if X is a fourfold, $\text{End}_K^0(X)$ is a CM-field of degree 4 and all endomorphisms of X are defined over K then there exists a finite algebraic extension L/K such that $\text{Isog}(X \times_K L, L)$ is infinite (see Theorem 10.3). A rather explicit example of the latter case (with infinite $\text{Isog}(X \times_K L, L)$) is provided by the field $K = \mathbf{C}(\lambda)$ of rational functions in independent variable λ , the jacobian X of the genus 4 curve $y^5 = x(x-1)(x-\lambda)$ and the overfield $L = \mathbf{C}(\sqrt[10]{\lambda}, \sqrt[10]{\lambda-1})$ (see Example 10.6).

The paper is organized as follows. Section 2 contains basic notation and useful facts about abelian varieties. In Section 3 we discuss abelian schemes over curves and corresponding monodromy representations. In Section 4 we state the main results. In Sections 5 and 6 we discuss non-isotrivial abelian schemes and isogenies of abelian schemes respectively. The next two sections contain the proofs of main results. Section 9 contains auxiliary results about quaternions. In Section 10 we deal with isogeny classes of four-dimensional abelian varieties.

Acknowledgements. This paper is based on beautiful ideas and results of J. Tate [32] and P. Deligne [6]. I learned about the Tate conjecture in the early 1970s from a paper of A. N. Parshin [24]. I am grateful to Minhyong Kim for stimulating questions and to B. van Geemen for useful comments. The final version of this paper was written during my stay at the Steklov Mathematical Institute of the Russian

Academy of Sciences (Moscow); I am grateful to its Department of Algebra for the hospitality.

2. ABELIAN VARIETIES

Let X be an abelian variety of positive dimension over a field K of characteristic zero. If n is a positive integer then we write X_n for the kernel of multiplication by n in $X(K_s)$. It is well-known [23] that X_n is a free $\mathbf{Z}/n\mathbf{Z}$ -module of rank $2\dim(X)$; it is also a Galois submodule in $X(\bar{K})$. We write $K(X_n)$ for the field of definition of all points of order n ; clearly, $K(X_n)/K$ is a finite Galois extension, whose Galois group is canonically identified with the image of $\text{Gal}(K)$ in $\text{Aut}(X_n)$. We write Id_X for the identity automorphism of X . We write $\text{End}(X)$ for the ring of all $\bar{K}$ -endomorphisms of X and $\text{End}^0(X)$ for the corresponding $\mathbf{Q}$ -algebra $\text{End}(X) \otimes \mathbf{Q}$. We have

$$\mathbf{Z} \cdot \text{Id}_X \subset \text{End}(X) \subset \text{End}^0(X).$$

Remark 2.1. Let X' be an abelian variety over K . If X and X' are K -isogenous then every K -isogeny $X \rightarrow X'$ gives rise to a bijection between $\text{Isog}(X, K)$ and $\text{Isog}(X', K)$. In particular, the sets $\text{Isog}(X, K)$ and $\text{Isog}(X', K)$ are either both finite or both infinite.

Let ℓ be a prime.

Lemma 2.2. *Suppose that ℓ is congruent to 1 modulo 4. If X admits a principal polarization over K and $\text{Isog}(X, K, \ell)$ is infinite then $\text{Isog}_1(X^2, K, \ell)$ is also infinite.*

Proof. Clearly, every abelian variety Y over K that admits an ℓ -isogeny $Y \rightarrow X$ over $\mathbf{C}(S)$ also admits a K -polarization, whose degree is a power of ℓ . In addition, there exists an ℓ -isogeny $X \rightarrow Y$ over K ; its dual $Y^t \rightarrow X^t \cong X$ is an ℓ -isogeny that is also defined over K . Since ℓ is congruent to 1 modulo 4 then $\sqrt{-1} \in \mathbf{Z}_\ell$ and it follows from Remark 5.3.1 on pp. 314–315 of [38] that $Y \times Y^t$ is principally polarized over K . Clearly, there is an ℓ -isogeny $Y \times Y^t \rightarrow X \times X = X^2$ that is defined over K . On the other hand, for any given abelian variety Z over K there are, up to an isomorphism, only finitely many abelian varieties over K that are isomorphic over K to an abelian subvariety of Z [16]. It follows that $\text{Isog}_1(X^2, K, \ell)$ is infinite. $\square$

Remark 2.3. Let L/K be a finite Galois extension with Galois group $\text{Gal}(L/K)$. Let Y be an abelian variety over K and $\text{Aut}_L(Y)$ be its group of L -automorphisms. It is well-known that $\text{Aut}_L(Y)$ is an *arithmetic* group; it follows from a theorem of Borel-Serre [1] that the corresponding first noncommutative Galois cohomology set $H^1(\text{Gal}(L/K), \text{Aut}_L(Y))$ is finite, i.e., the set of $(K$ -isomorphism classes of) L/K -forms of Y is finite. This implies that if $\text{Isog}(X \times_K L, L)$ (resp. $\text{Isog}(X \times_K L, L, \ell)$) is finite then $\text{Isog}(X, K)$ (resp. $\text{Isog}(X, K, \ell)$) is also finite [39].

Abusing notation, we sometime write $\text{Isog}(X, L, \ell)$, $\text{Isog}(X, L, \ell)$ and $\text{Isog}_m(X, L, \ell)$ instead of $\text{Isog}(X \times_K L, L, \ell)$, $\text{Isog}(X \times_K L, L, \ell)$ and $\text{Isog}_m(X \times_K L, L, \ell)$ respectively.

3. ABELIAN SCHEMES AND MONODROMY REPRESENTATIONS

Let S be an irreducible smooth (but not necessarily projective) algebraic curve over $\mathbf{C}$. We write $\mathbf{C}(S)$ for the field of rational functions on S and $\overline{\mathbf{C}(S)}$ for its algebraic closure. Let $f : \mathcal{X} \rightarrow S$ be a polarized abelian scheme of positive relative dimension d over S . Let η be the generic point of S and X be the generic fiber; it is a d -dimensional abelian variety over $k(\eta) = \mathbf{C}(S)$.

An abelian variety Z over $\mathbf{C}(S)$ is called *constant* if there exists an abelian variety W over $\mathbf{C}$ such that Z is isomorphic to $W \times_{\mathbf{C}} \mathbf{C}(S)$ over $\mathbf{C}(S)$.

Recall that $\mathcal{X}$ is called *isotrivial* if there exists a finite étale cover $S' \rightarrow S$ (with non-empty S') such that the pullback $\mathcal{X}_{S'} = \mathcal{X} \times_S S'$ is a constant abelian scheme over S' . We say that $\mathcal{X}$ is *weakly isotrivial* if (under the same assumptions on $S' \rightarrow S$) $\mathcal{X}_{S'}$ contains a non-zero constant abelian subscheme.

Theorem 3.1. (i) $\mathcal{X}$ is isotrivial if and only if there exists an abelian variety W over $\mathbf{C}$ such that the $\mathbf{C}(S)$ -abelian varieties X and $W \times_{\mathbf{C}} \mathbf{C}(S)$ are isomorphic over $\overline{\mathbf{C}(S)}$.
(ii) $\mathcal{X}$ is weakly isotrivial if and only if there exist an abelian variety W of positive dimension over $\mathbf{C}$ and an abelian subvariety $Z \subset X \times_{\mathbf{C}(S)} \overline{\mathbf{C}(S)}$ such that the $\overline{\mathbf{C}(S)}$ -abelian varieties $W \times_{\mathbf{C}} \overline{\mathbf{C}(S)}$ and Z are isomorphic.

We prove Theorem 3.1 in Section 5.

We write $R_1 f_* \mathbf{Z}$ for the corresponding local system of the first integral homology groups $H_1(\mathcal{X}_s, \mathbf{Z})$ of the fibres $\mathcal{X}_s = f^{-1}(s)$ on S . If $g : \mathcal{Y} \rightarrow S$ is another polarized abelian scheme over S with generic fiber f then the natural map

$$\mathrm{Hom}_S(\mathcal{Y}, \mathcal{X}) \rightarrow \mathrm{Hom}_{\mathbf{C}(S)}(Y, X)$$

is bijective. We write $R_1 f_* \mathbf{Q}$ for the corresponding local system $R_1 f_* \mathbf{Z} \otimes \mathbf{Q}$ of the first rational homology groups $H_1(\mathcal{X}_s, \mathbf{Q}) = H_1(\mathcal{X}_s, \mathbf{Z}) \otimes \mathbf{Q}$ of the fibres $\mathcal{X}_s = f^{-1}(s)$ on S and $\pi_1(S, s) \rightarrow \mathrm{Aut}(H_1(\mathcal{X}_s, \mathbf{Z})) \subset \mathrm{Aut}(H_1(\mathcal{X}_s, \mathbf{Q}))$ for the corresponding monodromy representation.

Remark 3.2. An abelian scheme $f : \mathcal{X} \rightarrow S$ is isotrivial if and only if the image of the monodromy representation $\pi_1(S, s) \rightarrow \mathrm{Aut}(H_1(\mathcal{X}_s, \mathbf{Z}))$ is finite [14] (see also [6, Sect. 4.1.3.3]).

3.3. Rigidity and specialization. It follows from the rigidity lemma [22, Ch. 6, Sect. 1, Cor. 6.2] that the natural homomorphism

$$\mathrm{Hom}_S(\mathcal{Y}, \mathcal{X}) \rightarrow \mathrm{Hom}(\mathcal{Y}_s, \mathcal{X}_s), \quad u \mapsto u_s$$

is injective. In particular, the natural ring homomorphism $\mathrm{End}_S(\mathcal{X}) \rightarrow \mathrm{End}(\mathcal{X}_s)$ is also injective. It is well-known that the natural homomorphisms

$$\mathrm{Hom}(\mathcal{Y}_s, \mathcal{X}_s) \rightarrow \mathrm{Hom}(H_1(\mathcal{Y}_s, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Z})), \quad \mathrm{End}(\mathcal{X}_s) \rightarrow \mathrm{End}(H_1(\mathcal{X}_s, \mathbf{Z}))$$

are embeddings. Taking the compositions, we get the embeddings

$$\mathrm{Hom}_S(\mathcal{Y}, \mathcal{X}) \rightarrow \mathrm{Hom}(H_1(\mathcal{Y}_s, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Z})), \quad \mathrm{End}_S(\mathcal{X}) \rightarrow \mathrm{End}(H_1(\mathcal{X}_s, \mathbf{Z}));$$

the images of $\mathrm{Hom}_S(\mathcal{Y}, \mathcal{X})$ and of $\mathrm{End}_S(\mathcal{X})$ lie in $\mathrm{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Y}_s, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Z}))$ and $\mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Z}))$ respectively. Further we will identify $\mathrm{End}_S(\mathcal{X})$ with its image in $\mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Z})) \subset \mathrm{End}(H_1(\mathcal{X}_s, \mathbf{Z}))$ and $\mathrm{Hom}_S(\mathcal{Y}, \mathcal{X})$ with its image in $\mathrm{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Y}_s, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Z}))$ respectively.

We have

$$\mathrm{Hom}_S^0(\mathcal{Y}, \mathcal{X}) := \mathrm{Hom}_S(\mathcal{Y}, \mathcal{X}) \otimes \mathbf{Q} \subset$$

$$\mathrm{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Y}_s, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Z})) \otimes \mathbf{Q} = \mathrm{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Y}_s, \mathbf{Q}), H_1(\mathcal{X}_s, \mathbf{Q})),$$

$$\mathrm{End}_S^0(\mathcal{X}) := \mathrm{End}_S(\mathcal{X}) \otimes \mathbf{Q} \subset \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Z})) \otimes \mathbf{Q} = \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Q})).$$

Notice that a theorem of Grothendieck [12, p. 60] (see also [6, Sect. 4.1.3.2]) implies that

$$\mathrm{Hom}_S(\mathcal{Y}, \mathcal{X}) = \mathrm{Hom}(\mathcal{Y}_s, \mathcal{X}_s) \bigcap \mathrm{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Y}_s, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Z})),$$

$$\mathrm{End}_S(\mathcal{X}) = \mathrm{End}(\mathcal{X}_s) \bigcap \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Z})).$$

It follows that

$$\mathrm{Hom}_S^0(\mathcal{Y}, \mathcal{X}) = \mathrm{Hom}^0(\mathcal{Y}_s, \mathcal{X}_s) \bigcap \mathrm{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Y}_s, \mathbf{Q}), H_1(\mathcal{X}_s, \mathbf{Q})).$$

Here $\mathrm{Hom}^0(\mathcal{Y}_s, \mathcal{X}_s) = \mathrm{Hom}(\mathcal{Y}_s, \mathcal{X}_s) \otimes \mathbf{Q} \subset \mathrm{Hom}_{\mathbf{Q}}(H_1(\mathcal{Y}_s, \mathbf{Q}), H_1(\mathcal{X}_s, \mathbf{Q}))$. In particular,

$$\mathrm{End}_S^0(\mathcal{X}) = \mathrm{End}^0(\mathcal{X}_s) \bigcap \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Q})).$$

3.4. Let Γ_s be the image of $\pi_1(S, s)$ in $\mathrm{Aut}(H_1(\mathcal{X}_s, \mathbf{Q}))$, let G_s is the Zariski closure of Γ_s in $\mathrm{GL}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}))$ and G_s^0 its identity component. By a theorem of Deligne [6, Cor. 4.2.9], G_s^0 is a *semisimple* algebraic $\mathbf{Q}$ -group. Recall [2, Ch. 1, Sect. 1.2] that G_s^0 has finite index in G_s . It follows that the intersection $\Gamma_s^0 := \Gamma_s \cap G_s^0$ is a normal subgroup of finite index in Γ_s and this index divides $[G_s : G_s^0]$.

The groups Γ_s and G_s do not depend on s in the following sense. Recall that S is arcwise connected with respect to the complex topology. If t is another point of S then every path γ in S from s to t defines an isomorphism $\gamma_* : H_1(\mathcal{X}_s, \mathbf{Z}) \cong H_1(\mathcal{X}_t, \mathbf{Z})$ such that $\Gamma_t = \gamma_* \Gamma_s \gamma_*^{-1}$ and therefore

$$G_t = \gamma_* G_s \gamma_*^{-1}, \quad G_t^0 = \gamma_* G_s^0 \gamma_*^{-1}.$$

Let $\mathrm{Hdg}(\mathcal{X}_s) \subset \mathrm{GL}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}))$ be the *Hodge group* of $\mathcal{X}_s$ [23] (see also [37, 18, 19]). Recall that $\mathrm{Hdg}(\mathcal{X}_s)$ is a connected reductive algebraic $\mathbf{Q}$ -group and its centralizer $\mathrm{End}_{\mathrm{Hdg}(\mathcal{X}_s)}(H_1(\mathcal{X}_s, \mathbf{Q}))$ in $\mathrm{End}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}))$ coincides with $\mathrm{End}^0(\mathcal{X}_s)$.

The following assertion is a special case of a theorem of Deligne [37, Th. 7.3] (see also [7, Prop. 7.5], [17]).

Theorem 3.5. *For all $s \in S$ outside a countable set, G_s^0 is a normal algebraic subgroup in $\mathrm{Hdg}(\mathcal{X}_s)$.*

Definition 3.6. We say that a point $s \in S$ is *in general position* (with respect to $\mathcal{X} \rightarrow S$) if G_s^0 is a normal algebraic subgroup in $\mathrm{Hdg}(\mathcal{X}_s)$. It follows from Theorem 3.5 that every s outside a countable set is in general position. In particular, a point in general position always does exist.

Corollary 3.7. *Suppose that s is in general position and G_s is connected. Then*

$$\mathrm{End}^0(\mathcal{X}_s) = \mathrm{End}_S^0(\mathcal{X}) = \mathrm{End}_{\mathbf{C}(S)}^0(X).$$

If, in addition, $\mathcal{X}$ is not isotrivial and $\mathrm{Hdg}(\mathcal{X}_s)$ is a $\mathbf{Q}$ -simple algebraic group then $\mathrm{End}_{\mathbf{C}(S)}^0(X) = \mathrm{End}_S^0(\mathcal{X}) = \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Q}))$.

Proof. We have $G_s = G_s^0 \subset \text{Hdg}(\mathcal{X}_s)$. Clearly,

$$\text{End}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}_{G_s}(H_1(\mathcal{X}_s, \mathbf{Q})) \supset \text{End}_{\text{Hdg}(\mathcal{X}_s)}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}^0(\mathcal{X}_s).$$

It follows that $\text{End}_S^0(\mathcal{X}) = \text{End}^0(\mathcal{X}_s) \cap \text{End}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}^0(\mathcal{X}_s)$. Assume now that $\mathcal{X}$ is not isotrivial and $\text{Hdg}(\mathcal{X}_s)$ is $\mathbf{Q}$ -simple. By Remark 3.2, Γ_s is infinite and therefore G_s has positive dimension. Since s is in general position, G_s is normal in $\text{Hdg}(\mathcal{X}_s)$. Now the simplicity of $\text{Hdg}(\mathcal{X}_s)$ implies that $\text{Hdg}(\mathcal{X}_s) = G_s$ and therefore

$$\text{End}^0(\mathcal{X}_s) = \text{End}_{G_s}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}_S^0(X) = \text{End}_{\mathbf{C}(S)}(X).$$

□

3.8. Base change. Let $S' \rightarrow S$ be a finite étale cover and $\mathcal{X}' = X \times_S S'$ the corresponding abelian S' -scheme. If a point $s' \in S'$ lies above $s \in S$ then (in obvious notations)

$$\mathcal{X}'_{s'} = \mathcal{X}_s, H_1(\mathcal{X}_s, \mathbf{Z}) = H_1(\mathcal{X}'_{s'}, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Q}) = H_1(\mathcal{X}'_{s'}, \mathbf{Q}), \text{Hdg}(\mathcal{X}'_{s'}) = \text{Hdg}(\mathcal{X}_s).$$

The fundamental group $\pi_1(S', s)$ of S' is a subgroup of finite index in $\pi_1(S, s)$ and therefore $\Gamma_{s'}$ is a subgroup of finite index in Γ_s . It follows that $G_{s'}$ is a subgroup of finite index in G_s ; in particular, $G_{s'}^0 = G_s^0$. This implies that s is in general position with respect to $\mathcal{X}$ if and only if s' is in general position with respect to $\mathcal{X}'$. It also follows that if G_s is connected then $G_{s'} = G_s$.

On the other hand, let $\Gamma^0 \subset \Gamma_s$ be a subgroup of finite index (e.g., $\Gamma^0 = \Gamma_s^0$). Clearly, its Zariski closure lies between G_s^0 and G_s ; in particular, if $\Gamma^0 \subset \Gamma_s^0$ then this closure coincides with G_s^0 . Let $\pi^0 \subset \pi_1(S, s)$ be the preimage of Γ^0 : it is a subgroup of finite index in $\pi_1(S, s)$. Let $S^0 \rightarrow S$ be a finite étale map of irreducible smooth algebraic curves attached to π^0 with $\pi_1(S^0) = \pi^0$. Clearly, the degree of $S^0 \rightarrow S$ coincides with the index $[\Gamma_s : \Gamma^0]$.

Notice that if Γ^0 is normal in Γ_s then $S^0 \rightarrow S$ is Galois with Galois group Γ_s/Γ^0 .

Fix a point $s_0 \in S^0$ that lies above s . Then the image of the monodromy representation $\pi_1(S^0, s_0) = \pi^0 \rightarrow \text{Aut}(H_1(\mathcal{X}_s, \mathbf{Q}))$ attached to the abelian S^0 -scheme $\mathcal{X} \times_S S^0$ coincides with Γ^0 . In other words, $\Gamma_{s_0} = \Gamma^0$ and therefore $G_{s_0} = G_s^0$ is connected if $\Gamma^0 \subset \Gamma_s^0$.

Example 3.9. Let $n \geq 3$ be an integer and $\Gamma^n \subset \Gamma_s$ the kernel of the reduction map modulo n

$$\Gamma_s \subset \text{Aut}(H_1(\mathcal{X}_s, \mathbf{Z})) \twoheadrightarrow \text{Aut}(H_1(\mathcal{X}_s, \mathbf{Z}/n\mathbf{Z})).$$

Here $H_1(\mathcal{X}_s, \mathbf{Z}/n\mathbf{Z}) = H_1(\mathcal{X}_s, \mathbf{Z}) \otimes \mathbf{Z}/n\mathbf{Z}$ is the first integral homology group of $\mathcal{X}_s$ with coefficients in $\mathbf{Z}/n\mathbf{Z}$. Clearly, Γ^n is a (normal) subgroup of finite index in Γ_s . On the other hand, since $\Gamma^n \subset 1 + n\text{End}(H_1(\mathcal{X}_s, \mathbf{Z}))$, Zariski closure of Γ^n is connected [31, Prop. 2.6]. This implies that this closure coincides with G_s^0 .

Remark 3.10. (i) It is known [30] that all endomorphisms of X are defined over $L := \mathbf{C}(S)(X_3)$. Clearly, $L/\mathbf{C}(S)$ is a finite Galois extension and all points of order 3 on X are L -rational. It follows from Néron-Ogg-Shafarevich criterion [29] that $L/\mathbf{C}(S)$ is unramified at all points of S . This implies that if S' is the normalization of S in K then $S' \rightarrow S$ is finite étale. Clearly, $\mathbf{C}(S') = L$.

- (ii) Suppose that G_s is connected. Then $\text{End}^0(X) = \text{End}_{\mathbf{C}(S)}^0(X)$, i.e., all endomorphisms of X are defined over $\mathbf{C}(S)$. Indeed, first,

$$\text{End}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}_{\Gamma_s}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}_{G_s}(H_1(\mathcal{X}_s, \mathbf{Q}))$$

and therefore $\text{End}_{\mathbf{C}(S)}^0(X) = \text{End}^0(\mathcal{X}_s) \cap \text{End}_{G_s}(H_1(\mathcal{X}_s, \mathbf{Q}))$. Second, pick a point $s' \in S'$ that lies above s . We have $\mathcal{X}'_{s'} = \mathcal{X}_s$, $G_s = G_{s'}$ and

$$\text{End}^0(X) = \text{End}_{\mathbf{C}(S')}^0(X) = \text{End}^0(\mathcal{X}'_{s'}) \cap \text{End}_{\pi_1(S',s')}(H_1(\mathcal{X}'_{s'}, \mathbf{Q})) =$$

$$\text{End}^0(\mathcal{X}'_{s'}) \cap \text{End}_{G_{s'}}(H_1(\mathcal{X}'_{s'}, \mathbf{Q})) = \text{End}^0(\mathcal{X}_s) \cap \text{End}_{G_s}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}_{\mathbf{C}(S)}^0(X).$$

3.11. Isogenies. It is well-known that $u \in \text{Hom}_{\mathbf{C}(S)}(Y, X)$ is an isogeny (resp. an ℓ -isogeny) if and only if there exist $v \in \text{Hom}_{\mathbf{C}(S)}(X, Y)$ and a positive integer n such that the compositions uv and vu are multiplications by n (resp. by ℓ^n) in X and Y respectively.

We say that $u \in \text{Hom}_S(\mathcal{Y}, \mathcal{X})$ is an isogeny (resp. an ℓ -isogeny) of abelian schemes if the induced homomorphism of generic fibers $u_\eta : Y \rightarrow X$ is an isogeny of abelian varieties.

Clearly, u is an isogeny (resp. an ℓ -isogeny) of abelian schemes if and only if there exist $v \in \text{Hom}_S(X, Y)$ and a positive integer n such that the compositions uv and vu are multiplications by n (resp. by ℓ^n) in $\mathcal{X}$ and $\mathcal{Y}$ respectively (see [4, Sect. 7.3]).

If $u \in \text{Hom}_S(\mathcal{Y}, \mathcal{X})$ is an isogeny (resp. an ℓ -isogeny) of abelian schemes then it is clear that the induced homomorphism $u_s : \mathcal{Y}_s \rightarrow \mathcal{X}_s$ is an isogeny (resp. an ℓ -isogeny) of the corresponding complex abelian varieties [4, Sect. 7.3].

Conversely, suppose that $u \in \text{Hom}_S(\mathcal{Y}, \mathcal{X})$ and assume that $u_s \in \text{Hom}(\mathcal{Y}_s, \mathcal{X}_s)$ is an isogeny (resp. an ℓ -isogeny). Then there exist an isogeny $w_s : \mathcal{X}_s \rightarrow \mathcal{Y}_s$ (resp. an ℓ -isogeny) and a positive integer n such that the compositions $u_s w_s$ and $w_s u_s$ are multiplications by n (resp. by ℓ^n) in $\mathcal{X}_s$ and $\mathcal{Y}_s$ respectively. Since $u_s \in \text{Hom}_{\pi_1(S,s)}(H_1(\mathcal{Y}_s, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Z}))$ and therefore

$$u_s^{-1} \in \text{Hom}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Z}) \otimes \mathbf{Q}, H_1(\mathcal{Y}_s, \mathbf{Z}) \otimes \mathbf{Q}),$$

it follows that $w_s \in \text{Hom}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Z}), H_1(\mathcal{Y}_s, \mathbf{Z}))$. This implies that

$$w_s \in \text{Hom}(\mathcal{X}_s, \mathcal{Y}_s) \cap \text{Hom}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Z}), H_1(\mathcal{Y}_s, \mathbf{Z}))$$

and therefore there exists $v \in \text{Hom}_S(\mathcal{X}, \mathcal{Y})$ with $v_s = w_s$. It follows that $(uv)_s$ and $(vu)_s$ are multiplications by n (resp. by ℓ^n) in $\mathcal{X}_s$ and $\mathcal{Y}_s$ respectively. By the rigidity lemma (Sect. 3.3), uv and vu are multiplications by n (resp. by ℓ^n) in $\mathcal{X}$ and $\mathcal{Y}$ respectively. This implies that u and v are isogenies (resp. ℓ -isogenies) of abelian schemes.

3.12. Semisimplicity. Recall [6, Sect. 4.2] that the *monodromy* representation

$$\pi_1(S, s) \rightarrow \text{Aut}(H_1(\mathcal{X}_s, \mathbf{Z}) \subset \text{Aut}(H_1(\mathcal{X}_s, \mathbf{Z}) \otimes \mathbf{Q})) = \text{Aut}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}))$$

is completely reducible and therefore its centralizer

$$D = D_f := \text{End}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Q})) \subset \text{End}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}))$$

is a finite-dimensional *semisimple* $\mathbf{Q}$ -algebra. In addition, the *center* E of D lies in $\text{End}_S(\mathcal{X}) \otimes \mathbf{Q}$ [6, Sect. 4.4.7]. It follows that E lies in the *center* C of $\text{End}_S^0(\mathcal{X})$.

Notice that $\text{End}_S(\mathcal{X}) \otimes \mathbf{Q} = \text{End}_{\mathbf{C}(S)}(X) \otimes \mathbf{Q}$. So, we have the inclusion of finite-dimensional semisimple $\mathbf{Q}$ -algebras

$$\text{End}_{\mathbf{C}(S)}(X) \otimes \mathbf{Q} =: \text{End}_S^0(\mathcal{X}) \subset D \subset \text{End}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}));$$

in addition, E lies in C . It follows easily that if D is commutative then $\text{End}_S^0(\mathcal{X}) = D$.

Remark 3.13. Assume that X is simple. Then:

- (i) $\text{End}_{\mathbf{C}(S)}^0(X)$ is a division algebra. Since the center E of D_f lies in $\text{End}_{\mathbf{C}(S)}^0(X)$ (Sect. 3.12), it has no zero divisors. This implies that E is a field and therefore D is a central simple E -algebra. It follows that the $H_1(\mathcal{X}_s, \mathbf{Q})$ is an isotypic $\pi_1(S, s)$ -module, i.e., is either simple or isomorphic to a direct sum of several copies of a simple module.
- (ii) Notice that $\text{End}_S^0(\mathcal{X}) = D$ if and only if $\text{End}_S(\mathcal{X}) = \text{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Z}))$. It is also known [6, Cor. 4.4.13] that $\text{End}_S^0(\mathcal{X}) = D$ when $d \leq 3$ and $\mathcal{X}$ is not weakly isotrivial.

Remark 3.14. Faltings [9, Sect. 5] has constructed a principally polarized abelian scheme $f : \mathcal{X} \rightarrow S$ with $d = 4$ that is not weakly isotrivial, $\Gamma_s = \Gamma^n$ for a certain integer $n \geq 3$ (in notations of Example 3.9) and $\text{End}_S^0(\mathcal{X}) \neq D$. It follows from arguments in Example 3.9 that in Faltings' example G_s is connected.

4. MAIN RESULTS

Our main result is the following statement.

Theorem 4.1. *Let $f : \mathcal{X} \rightarrow S$ be a polarized abelian scheme of positive relative dimension d . Then the following conditions are equivalent:*

- (i) $D_f \neq \text{End}_S^0(\mathcal{X})$.
- (ii) *The set $\text{Isog}(X, \mathbf{C}(S))$ is infinite.*

Theorem 4.1 is an immediate corollary of the following two statements.

Theorem 4.2. *Let $f : \mathcal{X} \rightarrow S$ be a polarized abelian scheme of positive relative dimension d . If $D_f = \text{End}_S^0(\mathcal{X})$ then $\text{Isog}(X, \mathbf{C}(S))$ is finite.*

Theorem 4.3. *Let $f : \mathcal{X} \rightarrow S$ be a polarized abelian scheme of positive relative dimension d . Suppose that $D = D_f \neq \text{End}_S^0(\mathcal{X})$ (in particular, D is noncommutative). If ℓ is a prime such that $D \otimes_{\mathbf{Q}} \mathbf{Q}_{\ell}$ is isomorphic to a direct sum of matrix algebras over fields then there exist a positive integer $d_0 < 2d$ and a sequence $\{Z(m)\}_{m=1}^{\infty}$ of $\mathbf{C}(S)$ -abelian varieties $Z(m)$ that enjoy the following properties:*

- (i) *For each positive integer m there exists a $\mathbf{C}(S)$ -isogeny $Z(m) \rightarrow X$ of degree $\ell^{(2d-d_0)m}$.*
- (ii) *Let Y be an abelian variety over $\mathbf{C}(S)$ and let M_Y be the set of positive integers m such that $Z(m)$ is isomorphic to Y over $\mathbf{C}(S)$. Then M_Y is either empty or finite. In other words, if M is an arbitrary infinite set of positive integers then there exists an infinite subset $M_0 \subset M$ such that for $m \in M_0$ all $Z(m)$ are mutually non-isomorphic over $\mathbf{C}(S)$.*

Remark 4.4. Clearly, $D \otimes_{\mathbf{Q}} \mathbf{Q}_{\ell}$ is isomorphic to a direct sum of matrix algebras over fields for all but finitely many primes ℓ .

We prove Theorems 4.3 and 4.2 in Sections 8 and 7 respectively.

Corollary 4.5. *Suppose that a point $s \in S$ is in general position and $\mathrm{Hdg}(\mathcal{X}_s)$ is $\mathbf{Q}$ -simple. If $\mathcal{X}$ is not weakly isotrivial then for all finite algebraic extension $L/\mathbf{C}(S)$ the set $\mathrm{Isog}(X, L)$ is finite.*

Proof. Replacing S by its normalization in L and using (3.8), we may assume without loss of generality that $L = K$. Choosing an integer $n \geq 3$, replacing S by S_n and applying (3.8) and Remark 2.3, we may assume without loss of generality that G_s is connected. It follows from Corollary 3.7 that

$$\mathrm{End}_{\mathbf{C}(S)}(X) = \mathrm{End}_S^0(\mathcal{X}) = \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Q})).$$

Now the result follows from Theorem 4.1. $\square$

Corollary 4.6. *Suppose that $D_f \neq \mathrm{End}_S^0(\mathcal{X})$. If X admits a principal polarization over $\mathbf{C}(S)$ then $\mathrm{Isog}_1(X^2, \mathbf{C}(S), \ell)$ is infinite for all but finitely many primes ℓ congruent to 1 modulo 4.*

Proof. Let us pick a prime ℓ such that $D_f \otimes_{\mathbf{Q}} \mathbf{Q}_{\ell}$ is a direct sum of matrix algebras over fields. It follows from Theorem 4.3 that $\mathrm{Isog}(X, \mathbf{C}(S), \ell)$ is infinite. Applying Lemma 2.2, we conclude that $\mathrm{Isog}_1(X^2, \mathbf{C}(S), \ell)$ is infinite if ℓ is congruent to 1 modulo 4. Now the result follows from Remark 4.4. $\square$

Till the end of this Section we assume that K is a field of algebraic functions in one variable over $\mathbf{C}$, i.e., K is finitely generated and of degree of transcendence 1 over $\mathbf{C}$.

Theorem 4.7. *Let X be an abelian variety over K of positive dimension d . Suppose that the $\bar{K}/\mathbf{C}$ -trace of X is zero. If $d \leq 3$ then $\mathrm{Isog}(X, K)$ is finite.*

Proof of Theorem 4.7. There exists a smooth irreducible algebraic $\mathbf{C}$ -curve S such that $K = \mathbf{C}(S)$ and there exists a polarized abelian scheme $\mathcal{X} \rightarrow S$ of relative dimension $d \leq 3$, whose generic fiber coincides with X [4, Sect.1.4, p.20]. Pick a point $s \in S(\mathbf{C})$. Recall [15, Ch. VIII, Sect. 3] that the vanishing of the $\overline{\mathbf{C}(S)}/\mathbf{C}$ -trace means that X does not contain over $\overline{\mathbf{C}(S)}$ a non-zero abelian subvariety that is isomorphic over $\overline{\mathbf{C}(S)}$ to a (constant) abelian variety of the form $W \times_{\mathbf{C}} \overline{\mathbf{C}(S)}$ where W is a complex abelian variety. By Theorem 3.1, this means X is not weakly isotrivial. It follows from results of Deligne (Remark 3.13(ii)) that $\mathrm{End}_S(\mathcal{X}) = \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Z}))$ and therefore $D_f := \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Q})) = \mathrm{End}_S^0(\mathcal{X})$. Now the assertion follows from Theorem 4.3. $\square$

Corollary 4.8. *Let X be an abelian variety over K that is isogenous over $\bar{K}$ to a product of abelian varieties of dimension ≤ 3 . If $\bar{K}/\mathbf{C}$ -trace of X is zero then $\mathrm{Isog}(X, K)$ is finite.*

Proof. Replacing if necessary, K by its finite Galois extension and using Remarks 2.3 and 2.1, we may assume without loss of generality that X is isomorphic over K to a product $Y_1 \times \cdots \times Y_r$ of abelian varieties Y_i 's of dimension ≤ 3 over K .

There exists a smooth irreducible algebraic $\mathbf{C}$ -curve S such that $K = \mathbf{C}(S)$ and there exist polarized abelian schemes $\mathcal{Y}_i \rightarrow S$, whose generic fiber coincides with Y_i [4, Sect.1.4, p.20]. If $f : \mathcal{X} \rightarrow S$ is the fiber product of $\mathcal{Y}_i$ then the generic fiber of the abelian S -scheme $\mathcal{X}$ coincides with X .

Pick a point $s \in S(\mathbf{C})$. Since all $\dim(Y_i) \leq 3$, it follows from results of Deligne [6, Cor. 4.4.13] applied to all pairs $(\mathcal{Y}_i, \mathcal{Y}_j)$ that $\mathrm{End}_S(\mathcal{X}) = \mathrm{End}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Z}))$

and therefore $D_f := \text{End}_{\pi_1(S,s)}(H_1(\mathcal{X}_s, \mathbf{Q})) = \text{End}_S^0(\mathcal{X})$. Now the assertion follows from Theorem 4.3. $\square$

Corollary 4.9. *Let X be a four-dimensional abelian variety over K . Suppose that the $\bar{K}/\mathbf{C}$ -trace of X is zero. If X is not absolutely simple then $\text{Isog}(X, K)$ is finite.*

Proof. Replacing if necessary, K by its finite Galois extension and using Remarks 2.3 and 2.1, we may assume without loss of generality that X is isomorphic over K to a product $Y \times T$ of abelian varieties Y and T of positive dimension over K . Since $4 = \dim(X) = \dim(Y) + \dim(T)$, we conclude that both $\dim(Y)$ and $\dim(T)$ do not exceed 3. One has only to apply Corollary 4.8. $\square$

Example 4.10. Let $f : \mathcal{X} \rightarrow S$ be the Faltings' example (Remark 3.14) and let X be its generic fiber, which is a principally polarized four-dimensional abelian variety over $\mathbf{C}(S)$. Since $\mathcal{X}$ is not weakly isotrivial, it follows from Theorem 3.1 that the $\overline{\mathbf{C}(S)}/\mathbf{C}$ -trace of X is zero. Since $D_f \neq \text{End}_S^0(\mathcal{X})$, it follows that $\text{Isog}(X, \mathbf{C}(S))$ is infinite and $\text{Isog}_1(X^2, \mathbf{C}(S), \ell)$ is also infinite for all but finitely many primes ℓ congruent to 1 modulo 4. Recall that G_s is connected (3.14) and therefore all endomorphisms of X are defined over $\mathbf{C}(S)$, thanks to Remark 3.10. It follows from Corollary 4.9 that X is absolutely simple.

We discuss isogeny classes of absolutely simple abelian fourfolds in Section 10.

5. NON-ISOTRIVIAL ABELIAN SCHEMES

Proof of Theorem 3.1. In one direction the assertion is almost obvious. Indeed, if $\mathcal{X}_{S'}$ is a constant abelian scheme (resp. contains a non-zero constant abelian subscheme) $W \times_{\mathbf{C}} S'$ where W is an abelian variety over $\mathbf{C}$ then $X \times_{\mathbf{C}(S)} \mathbf{C}(S')$ is isomorphic to $W \times_{\mathbf{C}} \mathbf{C}(S') = (W \times_{\mathbf{C}} \mathbf{C}(S)) \times_{\mathbf{C}(S)} \mathbf{C}(S')$ (resp. contains an abelian subvariety that is defined over $\mathbf{C}(S')$ and is isomorphic to $W \times_{\mathbf{C}} \mathbf{C}(S')$ over $\mathbf{C}(S')$). (By the way, we did not use an assumption that $S' \rightarrow S$ is étale.)

In the opposite direction, let W be an abelian variety over $\mathbf{C}$, let W_η be the constant abelian variety $W \times_{\mathbf{C}} \mathbf{C}(S)$ and $\bar{u} : W_\eta \times_{\mathbf{C}(S)} \overline{\mathbf{C}(S)} \hookrightarrow X \times_{\mathbf{C}(S)} \overline{\mathbf{C}(S)}$ an embedding of abelian varieties over $\overline{\mathbf{C}(S)}$. Let $L = \mathbf{C}(S)(X_3)$ be the field of definition of all points of order 3 on X and S' is the normalization of S in L . By Remark 3.10(i), $S' \rightarrow S$ is a finite étale map of smooth irreducible curves, $\mathbf{C}(S') = L$ and all points of order 3 on X are L -rational. Clearly, all torsion points of W_η are $\mathbf{C}(S)$ -rational; in particular, all points of order 3 on W_η are defined over L . It follows from results of [30] that all $\overline{\mathbf{C}(S)}$ -homomorphisms between W_η and X are defined over $\mathbf{C}(S')$; in particular, $\bar{u}$ is defined over $\mathbf{C}(S')$, i.e., there exists an embedding $u : W_\eta \times_{\mathbf{C}(S)} \mathbf{C}(S') \hookrightarrow X \times_{\mathbf{C}(S)} \mathbf{C}(S')$ of abelian varieties over $\mathbf{C}(S')$ such that $\bar{u}$ is obtained from u by extension of scalars from $\mathbf{C}(S')$ to $\overline{\mathbf{C}(S)}$. Notice that the $\mathbf{C}(S')$ -abelian varieties $X \times_{\mathbf{C}(S)} \mathbf{C}(S')$ and $W_\eta \times_{\mathbf{C}(S)} \mathbf{C}(S') = W \times_{\mathbf{C}} \mathbf{C}(S')$ are generic fibers of abelian S' -schemes $\mathcal{X}_{S'}$ and $W \times_{\mathbf{C}} S'$ respectively. It follows that u extends to a certain homomorphism of abelian S' -schemes $W \times_{\mathbf{C}} S' \rightarrow \mathcal{X}_{S'}$, which we denote by $u_{S'}$. If u is an isomorphism of generic fibers (the isotrivial case) then $u_{S'}$ is an isomorphism of the corresponding abelian schemes; in particular, $\mathcal{X}_{S'}$ is a constant abelian scheme. If u is not an isomorphism, it is still a closed embedding; in particular, the image Y of $u : W \times_{\mathbf{C}} \mathbf{C}(S') \hookrightarrow X \times_{\mathbf{C}(S)} \mathbf{C}(S')$ is an abelian subvariety in $X \times_{\mathbf{C}(S)} \mathbf{C}(S')$ and the abelian varieties Y and $W \times_{\mathbf{C}} \mathbf{C}(S')$ are isomorphic over $\mathbf{C}(S')$. Let $\mathcal{Y}$ be the schematic closure of the image Y in $\mathcal{X}_{S'}$.

It follows from Corollary 6 on p. 175 of [4] that $\mathcal{Y}$ is the Néron model of Y over S' . Since $W \times_{\mathbf{C}} \mathbf{C}(S') \cong Y$ over $\mathbf{C}(S')$ and the Néron model of $W \times_{\mathbf{C}} \mathbf{C}(S')$ over S' is $W \times S'$, we conclude that $\mathcal{X}_{S'}$ contains an abelian subscheme isomorphic to the constant abelian scheme $W \times_{\mathbf{C}} S'$. $\square$

6. ISOGENY CLASSES OF ABELIAN SCHEMES

Let $\mathcal{X} \rightarrow S$ be a polarized abelian scheme of positive relative dimension d . Let us consider the category $Is(\mathcal{X})$, whose objects are pairs $(\mathcal{Y}, \alpha)$ that consist of an abelian scheme $g : \mathcal{Y} \rightarrow S$ and an isogeny $\alpha : \mathcal{Y} \rightarrow \mathcal{X}$ of abelian schemes and the set of morphisms $\text{Mor}((\mathcal{Y}_1, \alpha_1), (\mathcal{Y}_2, \alpha_2)) := \text{Hom}_S(\mathcal{Y}_1, \mathcal{Y}_2)$ for any pair of objects $(\mathcal{Y}_1, \alpha_1)$ and $(\mathcal{Y}_2, \alpha_2)$. Let us consider the category $Is_s(\mathcal{X})$, whose objects are pairs (Λ, i) that consist of a $\pi_1(S, s)$ -module Λ , whose additive group is isomorphic to $\mathbf{Z}^{2d}$ and an embedding $i : \Lambda \hookrightarrow H_1(\mathcal{X}_s, \mathbf{Z})$ of $\pi_1(S, s)$ -modules and the set of morphisms $\text{Mor}_{Is_s(\mathcal{X})}((\Lambda_1, i_1), (\Lambda_2, i_2))$ is the set

$$\{a \in \text{Hom}_{\pi_1(S, s)}(\Lambda_1, \Lambda_2) \mid \exists u \in \text{End}^0(\mathcal{X}_s) \text{ such that } i_2 a = u i_1\}.$$

Theorem 6.1. *The functor*

$$\Psi_s : Is(\mathcal{X}) \rightarrow Is_s(\mathcal{X}), (\mathcal{Y}, \alpha) \mapsto (H_1(\mathcal{Y}_s, \mathbf{Z}), \alpha_s : H_1(\mathcal{Y}_s, \mathbf{Z}) \rightarrow H_1(\mathcal{X}_s, \mathbf{Z})),$$

$$(\gamma : \mathcal{Y}_1 \rightarrow \mathcal{Y}_2) \mapsto \gamma_s \in \text{Hom}(\mathcal{Y}_{1s}, \mathcal{Y}_{2s}) \bigcap \text{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Y}_{1s}, \mathbf{Z}), H_1(\mathcal{Y}_{2s}, \mathbf{Z}))$$

is an equivalence of categories.

Proof. First, we need to check that $\gamma_s \in \text{Mor}_{Is_s(\mathcal{X})}((H_1(\mathcal{Y}_{1s}, \mathbf{Z}), H_1(\mathcal{Y}_{2s}, \mathbf{Z})))$, i.e. there exists $u \in \text{End}^0(\mathcal{X}_s)$ such that $\alpha_{2s} \gamma_s = u \alpha_{1s}$. (Recall that $\alpha_1 : \mathcal{Y}_1 \rightarrow \mathcal{X}, \alpha_2 : \mathcal{Y}_2 \rightarrow \mathcal{X}$ are isogenies of abelian schemes and $\gamma_s \in \text{Hom}(\mathcal{Y}_{1s}, \mathcal{Y}_{2s})$.) Clearly, both $\alpha_{1s} : \mathcal{Y}_{1s} \rightarrow \mathcal{X}_s, \alpha_{2s} : \mathcal{Y}_{2s} \rightarrow \mathcal{X}_s$ are isogenies of abelian varieties. Then $u := \alpha_{2s} \gamma_s \alpha_{1s}^{-1} \in \text{End}^0(\mathcal{X}_s)$ satisfies $\alpha_{2s} \gamma_s = u \alpha_{1s}$. Second, the injectiveness and surjectiveness of $\text{Hom}_S(\mathcal{Y}_1, \mathcal{Y}_2) \rightarrow \text{Hom}(\mathcal{Y}_{1s}, \mathcal{Y}_{2s}) \bigcap \text{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Y}_{1s}, \mathbf{Z}), H_1(\mathcal{Y}_{2s}, \mathbf{Z}))$ follow from the rigidity lemma and Grothendieck's theorem (Sect. 3.3) respectively.

Let (Λ, i_s) be an object of $Is_s(\mathcal{X})$, i.e., a $\pi_1(S, s)$ -module Λ , whose additive group is isomorphic to $\mathbf{Z}^{2d}$ and an embedding $i_s : \Lambda \hookrightarrow H_1(\mathcal{X}_s, \mathbf{Z})$ of $\pi_1(S, s)$ -modules. In order to check the *essential surjectiveness* of Ψ_s , we need to construct an abelian scheme $g : \mathcal{Y} \rightarrow S$, an isogeny $\alpha : \mathcal{Y} \rightarrow \mathcal{X}$ and an isomorphism $\phi_s : \Lambda \cong H_1(\mathcal{Y}_s, \mathbf{Z})$ of $\pi_1(S, s)$ -modules such that $\phi_s = \alpha_s i_s$. In order to do that, recall that the $\pi_1(S, s)$ -module Λ defines a certain local system $\mathcal{U}$ of free $\mathbf{Z}$ -modules of rank $2d$ on S , whose fiber at s coincides with Λ . In addition, i_s defines an embedding of local systems $i : \mathcal{U} \hookrightarrow R_1 f_* \mathbf{Z}$, whose fiber at s coincides with our “original” i_s . Rank arguments imply that the corresponding embedding

$$i : \mathcal{U} \otimes \mathbf{Q} \rightarrow R_1 f_* \mathbf{Z} \otimes \mathbf{Q} = R_1 f_* \mathbf{Q}$$

is, in fact, an isomorphism. This allows us to provide $\mathcal{U}$ with the structure (induced by $R_1 f_* \mathbf{Z}$) of the holomorphic family of polarized Hodge structures of type $(-1, 0) + (0, -1)$ [6, Sect. 4.4]. The equivalence of the category of polarized abelian schemes and the category of holomorphic families of polarized Hodge structures of type $(-1, 0) + (0, -1)$ over S [6, Sect. 4.4.2 and 4.4.3] (based on results of [3]) implies that there exist an abelian scheme $g : \mathcal{Y} \rightarrow S$, a homomorphism $\alpha \in \text{Hom}_S(\mathcal{Y}, \mathcal{X})$ and an isomorphism of local systems $\phi_s : \mathcal{U} \cong R_1 g_* \mathbf{Z}$ such that $\alpha \psi_S = i$. Taking the fiber of the latter equality at s , we get the desired $\alpha_s (\psi_S)_s = i_s$. Clearly, $\alpha_s \in \text{Hom}(\mathcal{Y}_s, \mathcal{X}_s)$ induces an isomorphism $H_1(\mathcal{Y}_s, \mathbf{Q}) \cong H_1(\mathcal{X}_s, \mathbf{Q})$ and therefore

is an isogeny. Applying results of Section 3.11, we conclude that α is an isogeny of abelian schemes. $\square$

Remark 6.2. The degree of α coincides with the index $[H_1(\mathcal{X}_s, \mathbf{Z}) : i_s(\Lambda)]$.

7. FINITE ISOGENY CLASSES

Let L (resp. $L_{\mathbf{Q}}$) be the image of the group algebra $\mathbf{Z}[\pi_1(S, s)]$ (resp. of $\mathbf{Q}[\pi_1(S, s)]$) in $\text{End}(H_1(\mathcal{X}_s, \mathbf{Z}))$ (resp. in $\text{End}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}))$) induced by the monodromy representation. Clearly, L is an order in the $\mathbf{Q}$ -algebra $L_{\mathbf{Q}}$.

It follows from Jacobson's density theorem and the semisimplicity of the monodromy representation (over $\mathbf{Q}$) that $L_{\mathbf{Q}}$ is a semisimple $\mathbf{Q}$ -algebra that coincides with the centralizer $\text{End}_{D_f}(H_1(\mathcal{X}_s, \mathbf{Q}))$ of D_f and

$$\text{End}_{L_{\mathbf{Q}}}(H_1(\mathcal{X}_s, \mathbf{Q})) = D_f.$$

Clearly, each $\pi_1(S, s)$ -stable $\mathbf{Z}$ -lattice in $H_1(\mathcal{X}_s, \mathbf{Q})$ is an L -module, whose additive group is isomorphic to $\mathbf{Z}^{2d}$.

7.1. It follows from the Jordan-Zassenhaus theorem [27, Theorem 26.4] that there are, up to an isomorphism, only finitely many L -modules, whose additive group is isomorphic to $\mathbf{Z}^{2d}$.

Proof of Theorem 4.2. Since $D_f = \text{End}_{\mathbf{S}}^0(\mathcal{X})$, we have

$$D_f = \text{End}_{\mathbf{S}}^0(\mathcal{X}) \subset \text{End}^0(\mathcal{X}_s) \subset \text{End}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q})).$$

Thanks to Theorem 6.1, it suffices to check that the set of isomorphism classes of objects in $Is_s(\mathcal{X})$ is finite. Thanks to the Jordan-Zassenhaus theorem (Sect. 7.1), it becomes an immediate corollary of the following statement.

Lemma 7.2. *Suppose that (Λ_1, i_1) and (Λ_2, i_2) are objects in $Is_s(\mathcal{X})$.*

Then $i_1(\Lambda_1)$ and $i_2(\Lambda_2)$ are L -submodules in $H_1(\mathcal{X}_s, \mathbf{Z})$, whose additive group is isomorphic to $\mathbf{Z}^{2d}$. If the L -modules $i_1(\Lambda_1)$ and $i_2(\Lambda_2)$ are isomorphic then (Λ_1, i_1) and (Λ_2, i_2) are isomorphic.

Proof of Lemma 7.2. Recall that Λ_1 and Λ_2 are $\pi_1(S, s)$ -modules, whose additive groups are isomorphic to $\mathbf{Z}^{2d}$ and $i_1 : \Lambda_1 \hookrightarrow H_1(\mathcal{X}_s, \mathbf{Z})$, $i_2 : \Lambda_2 \hookrightarrow H_1(\mathcal{X}_s, \mathbf{Z})$ are embeddings of $\pi_1(S, s)$ -modules. Clearly, $i_1 : \Lambda_1 \cong i_1(\Lambda_1)$, $i_2 : \Lambda_2 \cong i_2(\Lambda_2)$ are isomorphisms of $\pi_1(S, s)$ -modules. Both $i_1(\Lambda_1)$ and $i_2(\Lambda_2)$ are L -submodules in $H_1(\mathcal{X}_s, \mathbf{Z})$,

Assume that there exists an isomorphism of L -modules $\alpha : i_1(\Lambda_1) \cong i_2(\Lambda_2)$. Clearly, α is an isomorphism of $\pi_1(S, s)$ -modules. Extending α by $\mathbf{Q}$ -linearity, we obtain an isomorphism

$$u : i_1(\Lambda_1) \otimes \mathbf{Q} \cong i_2(\Lambda_2) \otimes \mathbf{Q}$$

of $L \otimes \mathbf{Q}$ -modules. Recall that $L \otimes \mathbf{Q} = L_{\mathbf{Q}}$, $i_1(\Lambda_1) \otimes \mathbf{Q} = i_2(\Lambda_2) \otimes \mathbf{Q} = H_1(\mathcal{X}_s, \mathbf{Q})$. We have

$$u \in \text{End}_{L_{\mathbf{Q}}}(H_1(\mathcal{X}_s, \mathbf{Q})) \cap \text{Aut}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q})) = D_f \cap \text{Aut}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q})).$$

Since $D_f \subset \text{End}^0(\mathcal{X}_s)$, we obtain that $u \in \text{End}^0(\mathcal{X}_s)$. Since $u \in \text{End}^0(\mathcal{X}_s) \subset \text{End}_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}))$ is an automorphism of the finite-dimensional $\mathbf{Q}$ -vector space $H_1(\mathcal{X}_s, \mathbf{Q})$, we have $u^{-1} \in \text{End}^0(\mathcal{X}_s)$, i.e., $u \in \text{End}^0(\mathcal{X}_s)^*$. Now if we put

$$a := i_2^{-1} \alpha i_1 : \Lambda_1 \rightarrow i_1(\Lambda_1) \rightarrow i_2(\Lambda_2) \rightarrow \Lambda_2$$

then $a \in \text{Hom}_{\pi_1(S,s)}(\Lambda_1, \Lambda_2)$ is an isomorphism of $\pi_1(S,s)$ -modules and $i_2 a = u i_1$, $i_1 a^{-1} = u^{-1} i_2$. This implies that a is an isomorphism of (Λ_1, i_1) and (Λ_2, i_2) . $\square$

$\square$

8. INFINITE ISOGENY CLASSES

We deduce Theorem 4.3 from the two following auxiliary statements.

Lemma 8.1. *Let V be a finite-dimensional vector space over a field k , let Id_V be the identity automorphism of V , let $A \subset \text{End}_k(V)$ a k -subalgebra that contains $k \cdot \text{Id}_V$ and is isomorphic to a direct sum of matrix algebras over fields. Let $B \subset A$ be a semisimple k -subalgebra that contains the center of A but does not coincide with A .*

Then there exists a proper subspace $W \subset V$ that enjoys the following properties:

- (i) *There does exist $a \in A$ such that $a(V) = W$.*
- (ii) *There does not exist $b \in B$ such that $b(V) = W$.*

In order to state the next lemma, let us choose a prime ℓ and consider the natural $\mathbf{Q}_\ell$ -linear representation

$$\pi_1(S, s) \rightarrow \text{Aut}(H_1(\mathcal{X}_s, \mathbf{Q})) \subset \text{Aut}_{\mathbf{Q}_\ell}(H_1(\mathcal{X}_s, \mathbf{Q}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell) \subset \text{Aut}_{\mathbf{Q}_\ell}(H_1(\mathcal{X}_s, \mathbf{Q}_\ell));$$

here $H_1(\mathcal{X}_s, \mathbf{Q}_\ell) = H_1(\mathcal{X}_s, \mathbf{Q}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$. Clearly, this representation remains semisimple and the centralizer of $\pi_1(S, s)$ in $\text{End}_{\mathbf{Q}_\ell}(H_1(\mathcal{X}_s, \mathbf{Q}_\ell))$ coincides with $D_f \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$. Let us put $D = D_f$. We have

$$\text{End}_S^0(\mathcal{X}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell \subset D \otimes_{\mathbf{Q}} \mathbf{Q}_\ell \subset \text{End}_{\mathbf{Q}_\ell}(H_1(\mathcal{X}_s, \mathbf{Q}_\ell)).$$

Clearly, $E \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$ is the center of $D \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$ and lies in $C \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$, which is the center of $\text{End}_S^0(\mathcal{X}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$. It is also clear that both $\mathbf{Q}_\ell$ -algebras $D \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$ and $\text{End}_S^0(\mathcal{X}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$ are semisimple finite-dimensional.

Lemma 8.2. *Let ℓ be a prime, W a proper $\pi_1(S, s)$ -stable $\mathbf{Q}_\ell$ -vector space of $H_1(\mathcal{X}_s, \mathbf{Q}_\ell)$ and $d_0 = \dim_{\mathbf{Q}_\ell}(W)$. Then one of the following two conditions holds:*

- (1) *There exists $u \in D \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$ with $u(H_1(\mathcal{X}_s, \mathbf{Q}_\ell)) = W$.*
- (2) *There exists a sequence $\{Z(m)\}_{m=1}^\infty$ of $\mathbf{C}(S)$ -abelian varieties $Z(m)$ that enjoy the following properties:*
 - (i) *For each positive integer m there exists a $\mathbf{C}(S)$ -isogeny $Z(m) \rightarrow X$ of degree $\ell^{(2d-d_0)m}$.*
 - (ii) *Let Y be an abelian variety over $\mathbf{C}(S)$ and let M_Y be the set of positive integers m such that $Z(m)$ is isomorphic to Y over $\mathbf{C}(S)$. Then M_Y is either empty or finite.*

Remark 8.3. The statement (and the proof) of Lemma 8.2 is inspired by [32, Prop. 1, pp. 136–137].

Proof of Theorem 4.3(modulo Lemmas 8.1 and 8.2). Let us apply Lemma 8.1 to

$$k = \mathbf{Q}_\ell, V = H_1(\mathcal{X}_s, \mathbf{Q}_\ell), A = D \otimes_{\mathbf{Q}} \mathbf{Q}_\ell, B = \text{End}_S^0(\mathcal{X}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell.$$

We conclude that there exist a proper $\mathbf{Q}_\ell$ -vector subspace W in $H_1(\mathcal{X}_s, \mathbf{Q}_\ell)$, an element $v \in D \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$ with $v(H_1(\mathcal{X}_s, \mathbf{Q}_\ell)) = W$ but there does not exist an element $u \in \text{End}_S^0(\mathcal{X}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$ with $u(H_1(\mathcal{X}_s, \mathbf{Q}_\ell)) = W$. Since v lies in $D \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$, it commutes

with $\pi_1(S, s)$ and therefore W is $\pi_1(S, s)$ -stable. Now the result follows from Lemma 8.2. $\square$

Proof of Lemma 8.2. Let us consider the $\pi_1(S, s)$ -stable $\mathbf{Z}_\ell$ -lattice $H_1(\mathcal{X}_s, \mathbf{Z}_\ell) = H_1(\mathcal{X}_s, \mathbf{Z}) \otimes \mathbf{Z}_\ell$ in the $\mathbf{Q}_\ell$ -vector space $H_1(\mathcal{X}_s, \mathbf{Q}_\ell)$. Notice that for each positive integer m we have a canonical isomorphism of free $\mathbf{Z}/\ell^m \mathbf{Z}$ -modules

$$H_1(\mathcal{X}_s, \mathbf{Z}_\ell)/\ell^m H_1(\mathcal{X}_s, \mathbf{Z}_\ell) = H_1(\mathcal{X}_s, \mathbf{Z})/\ell^m H_1(\mathcal{X}_s, \mathbf{Z}) = H_1(\mathcal{X}_s, \mathbf{Z}/\ell^m \mathbf{Z})$$

(induced by the canonical isomorphism $\mathbf{Z}/\ell^m \mathbf{Z} = \mathbf{Z}_\ell/\ell^m \mathbf{Z}_\ell$) that commutes with the actions of $\pi_1(S, s)$.

The intersection $T := W \cap H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$ is a $\pi_1(S, s)$ -stable free pure $\mathbf{Z}_\ell$ -submodule of rank d_0 in $H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$. Clearly, $T \subset W$ and the natural map $T \otimes_{\mathbf{Z}_\ell} \mathbf{Q}_\ell \rightarrow W$ is an isomorphism of $\mathbf{Q}_\ell$ -vector spaces. The image T_m of T in $H_1(\mathcal{X}_s, \mathbf{Z}/\ell^m \mathbf{Z})$ is a free $\mathbf{Z}/\ell^m \mathbf{Z}$ -submodule of rank d_0 . The preimage of T_m in $H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$ coincides with $T + \ell^m H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$.

We write Λ_m for the preimage of T_m in $H_1(\mathcal{X}_s, \mathbf{Z})$. Clearly, $T + \ell^m H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$ contains Λ_m and the natural map $\Lambda_m \otimes \mathbf{Z}_\ell \rightarrow T + \ell^m H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$ is an isomorphism of free $\mathbf{Z}_\ell$ -modules. Notice that Λ_m is a $\pi_1(S, s)$ -stable subgroup of index $\ell^{m(2d-d_0)}$ in $H_1(\mathcal{X}_s, \mathbf{Z})$ and contains $\ell^m H_1(\mathcal{X}_s, \mathbf{Z})$. It follows from Theorem 6.1 that there exists an abelian scheme $h_m : \mathcal{Z}(m) \rightarrow S$ and an isogeny $\gamma(m) : \mathcal{Z}(m) \rightarrow \mathcal{X}$ of abelian schemes of degree $\ell^{m(2d-d_0)}$ such that $\gamma(m)_s(H_1(\mathcal{Z}(m)_s, \mathbf{Z})) = \Lambda_m$. (Here $\mathcal{Z}(m)_s$ is the fiber of $\mathcal{Z}(m)$ over s .) Since Λ_m contains $\ell^m H_1(\mathcal{X}_s, \mathbf{Z})$, there exists an isogeny $\gamma(m)'_s : \mathcal{X}_s \rightarrow \mathcal{Z}(m)_s$ of degree ℓ^{md_0} such that the compositions

$$\gamma(m)_s \gamma(m)'_s : \mathcal{X}_s \rightarrow \mathcal{Z}(m)_s \rightarrow \mathcal{X}_s, \quad \gamma(m)'_s \gamma(m)_s : \mathcal{Z}(m)_s \rightarrow \mathcal{X}_s \rightarrow \mathcal{Z}(m)_s$$

coincide with multiplication(s) by ℓ^m . Clearly,

$$\gamma(m)_s(H_1(\mathcal{Z}(m)_s, \mathbf{Z}_\ell)) = \gamma(m)_s(H_1(\mathcal{Z}(m)_s, \mathbf{Z})) \otimes \mathbf{Z}_\ell = \Lambda_m \otimes \mathbf{Z}_\ell = T + \ell^m H_1(\mathcal{X}_s, \mathbf{Z}_\ell).$$

We also have $H_1(\mathcal{Z}(m)_s, \mathbf{Z}) \supset \gamma(m)'_s(H_1(\mathcal{X}_s, \mathbf{Z})) \supset \ell^m H_1(\mathcal{Z}(m)_s, \mathbf{Z})$ and therefore

$$\begin{aligned} H_1(\mathcal{Z}(m)_s, \mathbf{Z}_\ell) \supset \gamma(m)'_s(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) \supset \gamma(m)'_s \gamma(m)_s(H_1(\mathcal{Z}(m)_s, \mathbf{Z}_\ell)) = \\ \ell^m H_1(\mathcal{Z}(m)_s, \mathbf{Z}_\ell). \end{aligned}$$

Since $\gamma(m)_s \in \text{Hom}_{\pi_1(S, s)}(H_1(\mathcal{Z}(m)_s, \mathbf{Z}), H_1(\mathcal{X}_s, \mathbf{Z}))$, we have

$$\gamma(m)_s^{-1} \in \text{Hom}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Q}), H_1(\mathcal{Z}(m)_s, \mathbf{Q})).$$

It follows that $\gamma(m)'_s = \ell^m \gamma(m)_s^{-1}$ lies in

$$\begin{aligned} \text{Hom}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Q}), H_1(\mathcal{Z}(m)_s, \mathbf{Q})) \cap \text{Hom}(H_1(\mathcal{X}_s, \mathbf{Z}), H_1(\mathcal{Z}(m)_s, \mathbf{Z})) = \\ \text{Hom}_{\pi_1(S, s)}(H_1(\mathcal{X}_s, \mathbf{Z}), H_1(\mathcal{Z}(m)_s, \mathbf{Z})). \end{aligned}$$

This implies that $\gamma(m)'_s$ coincides with the “fiber over” s of a certain isogeny of abelian schemes $\gamma(m)' : \mathcal{X} \rightarrow \mathcal{Z}(m)$ of degree ℓ^{md_0} . The rigidity lemma implies that the compositions

$$\gamma(m) \gamma(m)' : \mathcal{X} \rightarrow \mathcal{Z}(m) \rightarrow \mathcal{X}, \quad \gamma(m)' \gamma(m) : \mathcal{Z}(m) \rightarrow \mathcal{X} \rightarrow \mathcal{Z}(m)$$

coincide with multiplication(s) by ℓ^m .

The generic fiber $Z(m)$ of $\mathcal{Z}(m)$ is a $\mathbf{C}(S)$ -abelian variety and $\gamma(m)$ and $\gamma(m)'$ induce $\mathbf{C}(S)$ -isogenies $\gamma(m)_\eta : Z(m) \rightarrow X$, $\gamma(m)'_\eta : X \rightarrow Z(m)$. Their degrees are $\ell^{m(2d-d_0)}$ and ℓ^{md_0} respectively. Their composition(s)

$$\gamma(m)_\eta \gamma(m)'_\eta : X \rightarrow Z(m) \rightarrow X, \quad \gamma(m)'_\eta \gamma(m)_\eta : Z(m) \rightarrow X \rightarrow Z(m)$$

coincide with multiplication(s) by ℓ^m .

Suppose that the condition (2) is not fulfilled. Then there exist an infinite set I of positive integers and an abelian variety Y over $\mathbf{C}(S)$ such that Y and $Z(i)$ are isomorphic over $\mathbf{C}(S)$ for all $i \in I$. Let n be the smallest element of I . For each $i \in I$ let us fix a $\mathbf{C}(S)$ -isomorphism $v_{i,\eta} : Z(n) \rightarrow Z(i)$. Clearly, $v_{i,\eta}$ extends to an isomorphism of abelian schemes $\mathcal{Z}(n) \rightarrow \mathcal{Z}(i)$, which we denote by v_i . Since v_i is an isomorphism, its specialization $v_{i,s}$ at s satisfies

$$v_{i,s}(H_1(\mathcal{Z}(n)_s, \mathbf{Z}_\ell)) = H_1(\mathcal{Z}(i)_s, \mathbf{Z}_\ell).$$

Let us consider the composition

$$u_i := \gamma(i)v_i\gamma(n)' : \mathcal{X} \rightarrow \mathcal{Z}(n) \rightarrow \mathcal{Z}(i) \rightarrow \mathcal{X}.$$

We have

$$u_i \in \text{End}_S(\mathcal{X}) \subset \text{End}_S(\mathcal{X}) \otimes \mathbf{Z}_\ell \subset \text{End}_S(\mathcal{X}) \otimes \mathbf{Q}_\ell = \text{End}_S^0(\mathcal{X}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell.$$

Since

$$H_1(\mathcal{Z}(n)_s, \mathbf{Z}_\ell) \supset \gamma(n)'_s(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) \supset \ell^n H_1(\mathcal{Z}(n)_s, \mathbf{Z}_\ell),$$

we conclude that

$$H_1(\mathcal{Z}(i)_s, \mathbf{Z}_\ell) \supset v_i\gamma(n)'_s(H_1(\mathcal{Z}(i)_s, \mathbf{Z}_\ell)) \supset \ell^n H_1(\mathcal{Z}(i)_s, \mathbf{Z}_\ell)$$

and therefore

$$\ell^i H_1(\mathcal{X}_s, \mathbf{Z}_\ell) + T = \gamma(i)(H_1(\mathcal{Z}(i)_s, \mathbf{Z}_\ell)) \supset \gamma(i)v_i\gamma(n)'(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) \supset$$

$$\ell^n \gamma(i)(H_1(\mathcal{Z}(i)_s, \mathbf{Z}_\ell)) = \ell^n (\ell^i H_1(\mathcal{X}_s, \mathbf{Z}_\ell) + T) = \ell^{n+i} H_1(\mathcal{X}_s, \mathbf{Z}_\ell) + \ell^n T.$$

It follows that for all $i \in I$,

$$\ell^i H_1(\mathcal{X}_s, \mathbf{Z}_\ell) + T \supset \gamma(i)v_i\gamma(n)'(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) = u_i(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) \supset$$

$$\ell^{n+i} H_1(\mathcal{X}_s, \mathbf{Z}_\ell) + \ell^n T.$$

Recall that $\text{End}_S(\mathcal{X}) = \text{End}_{\mathbf{C}(S)}(X)$ is a free commutative group of finite rank and therefore the $\mathbf{Z}_\ell$ -lattice $\text{End}_S(\mathcal{X}) \otimes \mathbf{Z}_\ell$ in $\text{End}_S(\mathcal{X}) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$ is a compact metric space with respect to the ℓ -adic topology. So, we can extract from $\{u_i\}_{i \in I}$ a subsequence $\{u_j\}_{j \in J}$ that converges to a limit

$$u \in \text{End}_S(\mathcal{X}) \otimes \mathbf{Z}_\ell \subset \text{End}_{\mathbf{Z}_\ell}(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)).$$

We may assume that there is a sequence of nonnegative integers $\{m_j\}_{j \in J}$ that tends to infinity and such that for all j , $u - u_j \in \ell^{m_j} \cdot \text{End}_S(\mathcal{X}) \otimes \mathbf{Z}_\ell$. In particular, $(u - u_j)(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) \subset \ell^{m_j} \cdot H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$. It follows that

$$u(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) = \{\lim u_j(c) \mid c \in H_1(\mathcal{X}_s, \mathbf{Z}_\ell)\}.$$

This implies easily that $u(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) \subset T$. On the other hand, if $t \in \ell^n T$ then for each $j \in J$ there exists $c_j \in H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$ with $u_j(c_j) = t$. Since $H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$ is a compact metric space with respect to the ℓ -adic topology, we can extract from $\{c_j\}_{j \in J}$ a subsequence $\{c_k\}_{k \in K}$ that converges to a limit $c \in H_1(\mathcal{X}_s, \mathbf{Z}_\ell)$. Then $u(c) = \lim u_k(c) = \lim u_k(c_k) = t$. It follows that $\ell^n T \subset u(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) \subset T$. This implies that $u(H_1(\mathcal{X}_s, \mathbf{Q}_\ell)) = u(H_1(\mathcal{X}_s, \mathbf{Z}_\ell)) \otimes_{\mathbf{Z}_\ell} \mathbf{Q}_\ell = T \otimes_{\mathbf{Z}_\ell} \mathbf{Q}_\ell = W$. $\square$

Proof of Lemma 8.1. Step 1. Reduction to the case of simple A . Suppose that the semisimple k -algebra A splits into a direct sum $A = A_1 \oplus A_2$ of non-zero semisimple k -algebras A_1 and A_2 . Let e_1 and e_2 be the identity elements of A_1 and A_2 respectively. Clearly, $\text{Id}_V = e_1 + e_2, e_1 e_2 = e_2 e_1 = 0, e_1^2 = e_1, e_2^2 = e_2$. It is also clear that both e_1 and e_2 lie in the center of A and therefore in the center of B . Let us put

$$V_1 = e_1 V, V_2 = e_2 V, B_1 = e_1 B = B e_1 \subset A_1, B_2 = e_2 B = B e_2 \subset A_2;$$

we have $V = V_1 \oplus V_2, B = B_1 \oplus B_2$. In addition, A_1 acts trivially on V_2 and A_2 acts trivially on V_1 . So, we may view A_1 as a subalgebra of $\text{End}_k(V_1)$ and A_2 as a subalgebra of $\text{End}_k(V_2)$ respectively. Obviously, the center of A_i lies in the center of B_i for both $i = 1, 2$. Clearly, either $B_1 \neq A_1$ or $B_2 \neq A_2$. It is also clear that the validity of the assertion of Lemma 8.1 for (V_i, A_i, B_i) with $A_i \neq B_i$ implies its validity for (V, A, B) . It follows that it suffices to prove Lemma 8.1 under an additional assumption that A is simple. So, further we assume that A is a simple k -algebra and therefore is isomorphic to a matrix algebra $M_r(E)$ of size r over a field E . Since E is the center of $M_r(E)$, it is a finite algebraic extension of k . The field E lies in the center of D , i.e., D is a E -algebra. It follows easily that a semisimple k -algebra D is also a semisimple E -algebra.

Step 2. Reduction to the case of simple $V = E^r$. Since all $M_r(E)$ -modules of finite k -dimension are isomorphic to direct sums of finite number of copies of the standard module E^r , we may assume that $V = E^r, A = M_r(E) = \text{End}_E(V)$. Clearly, if W is k -vector subspace in V then one can find $a \in \text{End}(V)$ with $u(V) = W$ if and only if W is a E -vector subspace in V . So, in order to prove Lemma 8.1, it suffices to prove the following statement.

Proposition 8.4. *Let V be a vector space of finite dimension r over a field E . Let $B \subset \text{End}_E(V)$ be a semisimple E -subalgebra that contains Id_V . If $B \neq \text{End}_E(V)$ then there exists a proper E -vector subspace W of V that enjoys the following property: there does not exist $b \in B$ with $b(V) = W$.*

Proof of Proposition 8.4. Suppose that B is not simple, i.e. B splits into a direct sum $B = B_1 \oplus B_2$ of non-zero summands B_1 and B_2 . Let e_1 and e_2 be the identity elements of A_1 and A_2 respectively. Clearly V splits into a direct sum $V = e_1 V \oplus e_2 V$ of non-zero B -stable subspaces $e_1 V$ and $e_2 V$. It is also clear that for each $b \in B$ the image

$$bV = bV_1 \oplus bV_2, bV_1 \subset V_1, bV_2 \subset V_2.$$

In particular, if W is not a direct sum of a subspace of $e_1 V$ and a subspace of $e_2 V$ then it does not coincide with bV for any choice of $b \in B$. This proves Proposition 8.4 in the case of non-simple B . So further we assume that B is a simple E -algebra. Let F be the center of B . Clearly, F is a field that contains $E \cdot \text{Id}_V$. If $F \neq E \cdot \text{Id}_V$ then every $b(V)$ is an F -vector space; in particular, its k -dimension is divisible by the degree $[F : E \cdot \text{Id}_V]$, which is greater than 1. So, none of E -subspaces W of E -dimension 1 is of the form $b(V)$. So, without loss of generality, we may assume that $F = E \cdot \text{Id}_V$, i.e., B is a central simple F -algebra. Then there exist a division algebra H of finite dimension over its center E and a positive integer m such that B is isomorphic to the matrix algebra $M_m(H)$. It follows from the classification of modules over central simple algebras that the left $B = M_m(H)$ -module V is isomorphic to a direct sum of finitely many copies of the standard module H^m .

Clearly, V carries a natural structure of right H -module in such a way that every $b(V)$ is a right H -submodule. In particular, $\dim_E(bV)$ is divisible by $\dim_E(H)$. This implies that if $H \neq E$ then none of E -subspaces W of E -dimension 1 is of the form bV . So, without loss of generality, we may assume that $H = E$ and $B \cong M_m(E)$. Since $B \subset \text{End}_E(V)$ but $B \neq \text{End}_E(V)$, we conclude that $m < r$. Now dimension arguments imply that the $B = M_m(E)$ -module $V = E^r$ is isomorphic to the direct sum of r/m copies of the standard module E^m ; in particular, $m \mid r$. This implies that $\dim_E(bV)$ is divisible by r/m . Since $m < r$, none of E -subspaces W of E -dimension 1 is of the form bV . $\square$

$\square$

9. QUATERNIONS AND $\text{SU}(2)$

This stuff is (or should be) well-known. However, I was unable to find a proper reference. (However, see [26, Lecture 1, Example 12].)

Let E be a field of characteristic zero, F/E its quadratic extension. We write $\sigma : z \mapsto \bar{z}$ for the only nontrivial E -linear automorphism of F , which is an involution. Fix a non-zero element $\mathbf{i} \in E$ such that $\sigma(\mathbf{i}) = -\mathbf{i}$. Clearly,

$$F = E + E \cdot \mathbf{i}, \quad 0 \neq -a := \mathbf{i}^2 \in E.$$

Let V be a two-dimensional E -vector space. We write $\text{Aut}_F^1(V)$ for the group of F -linear automorphisms u of V with $\det_F(u) = 1$. Here $\det_F : \text{Aut}_F(V) \rightarrow F^*$ is the determinant map. If $\omega : V \times V \rightarrow F$ is a map then we write $\text{Aut}_F(V, \omega)$ for the group of all F -linear automorphisms u of V such that $\omega(ux, uy) = \omega(x, y) \forall x, y \in V$ and put

$$\text{Aut}_F^1(V, \omega) := \text{Aut}_F(V, \omega) \cap \text{Aut}_F^1(V).$$

Let $\psi : V \times V \rightarrow F$ be a non-degenerate F -sesquilinear Hermitian form. Let us reduce H to a diagonal form, i.e., pick $e_1 \in V$ with $\psi(e_1, e_1) \neq 0$ and let V_2 be the orthogonal complement in V to e_1 with respect to ψ . The non-degeneracy of ψ implies that V_2 is one-dimensional and the restriction of ψ to V_2 is also non-degenerate, i.e., ψ does not vanish on non-zero elements of V_2 . It follows that if e_2 is a non-zero element of V_2 then $\{e_1, e_2\}$ is an orthogonal basis of V then

$$\psi(z_1 e_1 + z_2 e_2, w_1 e_1 + w_2 e_2) = b_1 z_1 \bar{w}_1 + b_2 z_2 \bar{w}_2 \quad \forall z_1, z_2, w_1, w_2 \in F$$

where $0 \neq b_1 := \psi(e_1, e_1) \in E$, $0 \neq b_2 := \psi(e_2, e_2) \in E$. Let us put $\text{discr}(\psi) = b_1 b_2$. It is known [5, Ch. 9, §2] that the (multiplicative) class of $\text{discr}(\psi)$ in $E^*/N_{F/E}(F^*)$ does not depend on the choice of basis. Here $N_{F/E} : F \rightarrow E$ is the norm map. Let us put $b = b_2/b_1 \in F^*$ and consider the Hermitian form

$$\psi' := \frac{1}{b_1} \psi : V \times V \rightarrow F,$$

$$\psi'(z_1 e_1 + z_2 e_2, w_1 e_1 + w_2 e_2) = z_1 \bar{w}_1 + b z_2 \bar{w}_2 \quad \forall z_1, z_2, w_1, w_2 \in F.$$

Clearly, the classes of b and $\text{discr}(\psi)$ in $E^*/N_{F/E}(F^*)$ do coincide and

$$\text{Aut}_F^1(V, \psi') = \text{Aut}_F^1(V, \psi).$$

Let us consider the cyclic algebra [25, Sect. 15.1]

$$D = (F, \sigma, -b) \cong (F, \sigma, -\text{discr}(\psi)).$$

Recall that D is the four-dimensional central simple E -algebra that contains F and coincides as (left) F -vector space with $F \oplus F \cdot \mathbf{j}$ where $\mathbf{j}$ is an element of A^* such that $\mathbf{j}^2 = -b$, $\mathbf{j}z\mathbf{j}^{-1} = \sigma(z) = \bar{z} \forall z \in F$. If we put $\mathbf{k} := \mathbf{i}\mathbf{j}$ then

$$A = F \oplus F \cdot \mathbf{j} = E \cdot 1 \oplus E \cdot \mathbf{i} \oplus E\mathbf{j} \oplus E \cdot \mathbf{k}$$

with

$$\mathbf{i}^2 = -a, \mathbf{j}^2 = -b, \mathbf{k}^2 = -ab, \mathbf{k} = \mathbf{i}\mathbf{j} = -\mathbf{j}\mathbf{i}, \mathbf{i}\mathbf{k} = -a\mathbf{j} = -\mathbf{k}\mathbf{i}, \mathbf{k}\mathbf{j} = -b\mathbf{i} = -\mathbf{j}\mathbf{k}.$$

Clearly, $D = \left(\frac{-a, -b}{E}\right)$ (see [25, Sect. 15.4]). Let us consider the standard F -linear involution $D \rightarrow D$, $q \mapsto \bar{q}$ that sends $\mathbf{i}, \mathbf{j}, \mathbf{k}$ to $-\mathbf{i}, -\mathbf{j}, -\mathbf{k}$ respectively. Clearly,

$$\bar{q}q = q\bar{q} = x^2 + ay^2 + bs^2 + abt^2 \subset E \forall q = x \cdot 1 + y \cdot \mathbf{i} + s \cdot \mathbf{j} + t \cdot \mathbf{k}; \quad x, y, s, t \in E.$$

Let us consider the “quaternionic” Hermitian D -sesquilinear form

$$\phi_D : D \times D \rightarrow D, (q_1, q_2) \mapsto q_1 \bar{q}_2.$$

Taking the compositions of $\phi_D : D \times D \rightarrow D$ with the projection maps

$$D = F \oplus F \cdot \mathbf{j} \twoheadrightarrow F, \quad D = F \oplus F \cdot \mathbf{j} \twoheadrightarrow F\mathbf{j},$$

we get E -bilinear forms

$$H_D : D \times D \rightarrow F, \quad A_D : D \times D \rightarrow F$$

defined by

$$\phi_D(q_1, q_2) = H_D(q_1, q_2) \cdot 1 + A_D(q_1, q_2) \cdot \mathbf{j}.$$

Clearly, both H_D and A_D are F -linear with respect to first argument. Since

$$\phi_D(q_1, q_2) = \overline{\phi_D(q_2, q_1)}, \quad \phi_D(q, q) = q\bar{q} \in E \subset F,$$

we have

$$H_D(q_1, q_2) = \overline{H_D(q_2, q_1)}, \quad A_D(q, q) = 0 \quad \forall q_1, q_2, q \in D.$$

This means that H_D is an Hermitian F -sesquilinear form on D and A_D is an alternating F -bilinear form on D .

Clearly, A_D is not identically zero and therefore is non-degenerate, since the F -dimension of D is 2. This implies that the *symplectic* group $\text{Aut}_F(D, A_D)$ coincides with $\text{Aut}_F^1(V)$ and therefore

$$\begin{aligned} \text{Aut}_F(D, \phi_D) &= \text{Aut}_F(D, H_D) \bigcap \text{Aut}_F(D, A_D) = \text{Aut}_F(D, H_D) \bigcap \text{Aut}_F^1(V) = \\ &= \text{Aut}_F^1(D, H_D). \end{aligned}$$

Clearly, $F \cdot 1$ and $F \cdot \mathbf{j}$ are mutually orthogonal with respect to H_D . On the other hand

$$H_D(z_1 \cdot 1, z_2 \cdot 1) = z_1 \bar{z}_2, \quad H_D(z_1 \cdot \mathbf{j}, z_2 \cdot \mathbf{j}) = z_1 \mathbf{j} \bar{z}_2 \mathbf{j} = z_1 \mathbf{j} (-\mathbf{j}) \bar{z}_2 = b z_1 \bar{z}_2 \quad \forall z_1, z_2 \in F.$$

This implies that the isomorphism of F -vector spaces

$$\kappa : V \cong D, \quad ze_1 + we_2 \mapsto z \cdot 1 + w \cdot \mathbf{j}$$

is an isomorphism of Hermitian F -vector spaces (V, ψ') and (D, H_D) , i.e.,

$$H_D(\kappa(v_1), \kappa(v_2)) = \psi'(v_1, v_2) \quad \forall v_1, v_2 \in V.$$

For every $q \in D$ we denote by $R(q)$ the F -linear operator $D \rightarrow D$, $d \mapsto d \cdot q$. Clearly,

$$R(1) = \text{Id}_D, \quad R(q_1 q_2) = R(q_2) R(q_1), \quad R(x q_1 + y q_2) = x R(q_1) + y R(q_2)$$

for all $q_1, q_2 \in D$; $x, y \in E$. I claim that

$$\text{Aut}_F(D, \phi_D) = \{R(q) \mid q\bar{q} = 1\}.$$

Indeed, if $q\bar{q} = 1$ then

$$\phi_D(R(q)q_1, R(q)q_2) = \phi_D(q_1q, q_2q) = q_1q\overline{q_2q} = q_1q\bar{q}\overline{q_2} = q_1\overline{q_2} = \phi_D(q_1, q_2),$$

i.e., $R(q)$ preserves ϕ_D . On the other hand, if a E -linear automorphism u of D preserves ϕ_D then $u(1)\overline{u(1)} = \phi_D(u(1), u(1)) = \phi_D(1, 1) = 1\bar{1} = 1 \cdot 1 = 1$ and $u' = R(u(1))^{-1}u$ also preserves ϕ_D and satisfies $u'(1) = 1$. This implies that for all $q \in D$,

$$q = q \cdot 1 = q = q \cdot \bar{1} = \phi_D(q, 1) = \phi_D(u'(q), u'(1)) = u'(q) \cdot \bar{1} = u'(q),$$

i.e., u' is the identity map Id_D and therefore $u = R(u(1))$. Since $\text{Aut}_F(D, \phi_D) = \text{Aut}_F^1(D, H_D)$, we conclude that $\text{Aut}_F^1(D, H_D) = \{R(q) \mid q\bar{q} = 1\}$.

Viewing D as the left D -module, we get an embedding $D \subset \text{End}_E(D)$. Clearly, (the algebra) D (of left multiplications) commutes with all right multiplications $R(q)$ and therefore with $\text{Aut}_F^1(D, H_D)$ in $\text{End}_E(D)$.

Lemma 9.1. *The centralizer $\mathcal{D}$ of $\text{Aut}_F^1(D, H_D)$ in $\text{End}_E(D)$ coincides with D .*

Proof. Clearly, $\mathcal{D}$ contains D . Let us pick non-zero integers n and m such that $n^2 + a \neq 0$, $m^2 + b \neq 0$ and put

$$q_1 = \frac{(n^2 - a) \cdot 1 + 2n \cdot \mathbf{i}}{n^2 + a}, \quad q_2 = \frac{(m^2 - b) \cdot 1 + 2m \cdot \mathbf{j}}{m^2 + b}.$$

One may easily check that $q_1\bar{q}_1 = 1 = q_2\bar{q}_2$. Clearly,

$$R(q_1) = \frac{n^2 - a}{n^2 + a} \text{Id}_V + \frac{2n}{n^2 + a} R(\mathbf{i}), \quad R(q_2) = \frac{m^2 - b}{m^2 + b} \text{Id}_V + \frac{2m}{m^2 + b} R(\mathbf{j})$$

and therefore $\mathcal{D}$ commutes with $R(q_1)$ and $R(q_2)$. Since n and m do not vanish, we conclude that $\mathcal{D}$ commutes with right multiplications $R(\mathbf{i})$ and $R(\mathbf{j})$. Clearly, $R(\mathbf{j})R(\mathbf{i}) = R(\mathbf{k})$. This implies that $\mathcal{D}$ commutes with $R(\mathbf{k})$ and therefore commutes with all $R(q)$ ($q \in D$). Now if $u \in \mathcal{D}$ then let us put $z := u(1) \in D$. For all $q \in D$ we have $u(q) = u(R(q)(1)) = R(q)(u(1)) = R(q)(z) = zq$, i.e., $u = z \in D \subset \text{End}_E(D)$. $\square$

Recall that the Hermitian F -vector spaces (V, ϕ') and (D, H_D) are isomorphic. Taking into account that $\text{Aut}_F^1(V, \psi) = \text{Aut}_F^1(V, \psi')$ and applying Lemma 9.1, we obtain the following statement.

Theorem 9.2. *Let us view V as a four-dimensional E -vector space. Then the centralizer D of $\text{Aut}_F^1(V, \psi)$ in $\text{End}_E(V)$ is a central simple four-dimensional E -algebra isomorphic to the four-dimensional central simple E -algebra $D = \left(\frac{-a, -b}{E}\right)$.*

10. ABELIAN FOURFOLDS

Theorem 10.1. *Let S be a smooth irreducible algebraic curve over $\mathbf{C}$ and $f : \mathcal{X} \rightarrow S$ a polarized abelian scheme of relative dimension 4 with generic fiber X .*

Suppose that $\mathcal{X}$ is not weakly isotrivial and $\text{Isog}(X, \mathbf{C}(S))$ is infinite. Then:

- (i) X is absolutely simple.
- (ii) The center E of D_f is a real quadratic field.

- (iii) D_f is a quaternion division E -algebra that is unramified at one infinite place of E and ramified at the other infinite place.
- (iv) $\text{End}_{\mathbf{C}(S)}^0(X)$ is either E or a CM-field of degree 4.
- (v) $\text{End}^0(X)$ is a CM-field of degree 4.
- (vi) Let $s \in S$ and assume that G_s is connected. Then $\text{End}_{\mathbf{C}(S)}^0(X) = \text{End}^0(X)$ is a CM-field of degree 4.

Proof. The absolute simplicity of X follows from Corollary 4.9, which proves (i). This implies that $\text{End}_{\mathbf{C}(S)}^0(X)$ has no zero divisors. Since E is isomorphic to a subalgebra of $\text{End}_{\mathbf{C}(S)}^0(X)$, we conclude that E is a number field and $[E : \mathbf{Q}]$ divides $2\dim(X) = 8$. Since $\text{Isog}(X, \mathbf{C}(S))$ is infinite, it follows from Theorem 4.1 that $D_f \neq \text{End}_S^0(\mathcal{X})$. By the last sentence of 3.12, $D_f \neq E$, i.e., D_f is a non-commutative central simple E -algebra. Since $\dim_E(D_f)$ divides

$$\dim_E(H_1(\mathcal{X}_s, \mathbf{Q})) = \frac{\dim_{\mathbf{Q}}(H_1(\mathcal{X}_s, \mathbf{Q}))}{[E : \mathbf{Q}]} = \frac{8}{[E : \mathbf{Q}]},$$

we conclude that $8/[E : \mathbf{Q}]$ is *not* square-free. It follows that E is either $\mathbf{Q}$ or a quadratic field. On the other hand, Deligne [6, Prop. 4.4.11] proved that if E is either $\mathbf{Q}$ or an imaginary quadratic field then $D_f = \text{End}_S^0(\mathcal{X})$. It follows that E is a real quadratic field (which proves (ii)) and $\dim_E(D_f) = 4$. It follows from [6, Prop. 4.4.11] combined with the inequality $D_f \neq \text{End}_S^0(\mathcal{X})$ that D_f is unramified at one infinite place of E and ramified at the another one. This rules out the possibility that D_f is a matrix algebra over E . It follows that D_f is a quaternion division E -algebra, which proves (iii).

We have

$$E \subset \text{End}_S^0(\mathcal{X}) \subset D_f, \quad \text{End}_S^0(\mathcal{X}) \neq D_f.$$

Since $\dim_E(D_f) = 4$, we conclude that either $\text{End}_S^0(\mathcal{X}) = E$ or $\text{End}_S^0(\mathcal{X})$ is a field of degree 4 that contains E . Since there is an embedding $E \hookrightarrow \mathbf{R}$ such that $D_f \otimes_E \mathbf{R}$ is the standard quaternion $\mathbf{R}$ -algebra, either $E = \text{End}_S^0(\mathcal{X})$ or $\text{End}_S^0(\mathcal{X})$ is a degree four field that is *not* totally real. Since $\text{End}_S^0(\mathcal{X}) = \text{End}_{\mathbf{C}(S)}^0(X)$, we conclude that either $E = \text{End}_S^0(\mathcal{X}) = \text{End}_{\mathbf{C}(S)}^0(X)$ or $\text{End}_S^0(\mathcal{X}) = \text{End}_{\mathbf{C}(S)}^0(X)$ is a CM-field of degree 4, which proves (iv).

Let $s \in S$ be a point in general position and assume that G_s is connected. Suppose that $\text{End}_{\mathbf{C}(S)}^0(X) = E$. We need to arrive to a contradiction. It follows from the first assertion of Corollary 3.7 that

$$\text{End}^0(\mathcal{X}_s) = \text{End}_S^0(\mathcal{X}) = \text{End}_{\mathbf{C}(S)}^0(X).$$

This implies that $\text{End}^0(\mathcal{X}_s) = E$ is a real quadratic field. It follows from [18, 4.2, p. 566] that $\text{Hdg}(\mathcal{X}_s)$ is $\mathbf{Q}$ -simple. It follows from the first assertion of Corollary 3.7 that $\text{End}_{\mathbf{C}(S)}^0(X) = \text{End}_S^0(X) = \text{End}_{\pi_1(S, s)}^0(H_1(\mathcal{X}_s, \mathbf{Q}))$. Applying Theorem 4.1, we conclude that $\text{Isog}(X, K)$ is finite. The obtained contradiction proves that $\text{End}_{\mathbf{C}(S)}^0(X)$ is a CM-field of degree 4. It follows from Remark 3.10 that $\text{End}_{\mathbf{C}(S)}^0(X) = \text{End}^0(X)$. This proves (vi).

We still have to prove that $\text{End}^0(X)$ is a CM-field of degree 4 without assuming the connectedness of G_s . However, there exists a (connected) finite étale cover $S' \rightarrow S$ such that for $s' \in S'$ the group $G_{s'}$ attached to the abelian scheme $\mathcal{X}' = \mathcal{X} \times_S S'$ is connected and all endomorphisms of X are defined over the field $L = \mathbf{C}(S')$ (see Subsect. 3.8, Example 3.9 and Remark 3.10). By Remark 2.3, $\text{Isog}(X \times_K L, L)$ is

also infinite. Applying already proven (vi) to the generic fiber $X \times_K L$ of $\mathcal{X}'$, we conclude that $\text{End}^0(X) = \text{End}^0(X \times_K L)$ is a CM-field of degree 4. This proves (v). $\square$

Example 10.2. Let $f : \mathcal{X} \rightarrow S$ be the Faltings' example (Sect. 3.14) and 4.10) and let X be its generic fiber. It follows from Theorem 10.1 combined with arguments in Example 4.10 that $\text{End}_{\mathbf{C}(S)}^0(X) = \text{End}^0(X)$ is a CM-field of degree 4 that is a purely imaginary quadratic extension of a totally real quadratic field E and D_f is a quaternion division E -algebra that is unramified at one infinite place of E and ramified at the other infinite place. In fact, the quaternion algebra D_f was the starting point of Faltings' construction [9, Sect. 5].

Theorem 10.3. *Let S be a smooth irreducible algebraic curve over $\mathbf{C}$ and $f : \mathcal{X} \rightarrow S$ a polarized abelian scheme of relative dimension 4 with generic fiber X . Suppose that $\mathcal{X}$ is not isotrivial and $\text{End}^0(X) = \text{End}_{\mathbf{C}(S)}^0(X)$. Assume, in addition, that $F := \text{End}^0(X)$ is a CM-field of degree 4. Let μ_F be the (multiplicative) group of roots of unity in F and r_F its order. Then:*

- (0) $\mathcal{X}$ is not weakly isotrivial.
- (i) *There exists a finite étale cover $S^0 \rightarrow S$ such that $\mathbf{C}(S')/\mathbf{C}(S)$ is a cyclic extension of degree dividing r_F and the set $\text{Isog}(X, \mathbf{C}(S^0))$ is infinite.*
- (ii) *Let $s \in S$ and assume that G_s is connected. Then $\text{Isog}(X, \mathbf{C}(S))$ is infinite.*

Proof. Clearly, X is an absolutely simple abelian variety. It follows easily that $\mathcal{X}$ is not weakly isotrivial.

Let us put $F := \text{End}_{\mathbf{C}(S)}^0(X)$. By assumption, it is a purely imaginary quadratic extension of a real quadratic field E . Pick a point s in general position. By Corollary 3.7, $\text{End}^0(\mathcal{X}_s) = F$. Let us put $V := H_1(\mathcal{X}_s, \mathbf{Q})$; recall that V is an 8-dimensional $\mathbf{Q}$ -vector space. Clearly, V carries a natural structure of two-dimensional F -vector space and $\Gamma_s \subset \text{Aut}_F(V)$. We can do better, using the polarization on $\mathcal{X}$ that induces a polarization on $\mathcal{X}_s$, whose Riemann form gives rise to a non-degenerate alternating $\pi_1(S, s)$ -invariant $\mathbf{Q}$ -bilinear form $\phi : V \times V \rightarrow \mathbf{Q}$. Since the complex conjugation $e \mapsto e'$ on F is the only positive involution on the CM-field $F = \text{End}^0(\mathcal{X}_s)$, all Rosati involutions on $\text{End}^0(\mathcal{X}_s) = F$ coincide with the complex conjugation. This implies that $\phi(ex, y) = \phi(x, e'y) \forall x, y \in V, e \in F$ where $e \mapsto e'$ is the complex conjugation on F . Pick a non-zero element $\alpha \in E$ with $\alpha' = -\alpha$. Then there exists a unique (non-degenerate) F -Hermitian form

$$\psi : V \times V \rightarrow F$$

such that $\phi(x, y) = \text{tr}_{F/\mathbf{Q}}(\alpha\psi(x, y)) \forall x, y \in V$ [8, Sect. 9]. Here $\text{tr}_{F/\mathbf{Q}} : F \rightarrow \mathbf{Q}$ is the trace map. Since ϕ is $\pi_1(S, s)$ -invariant, ψ is also $\pi_1(S, s)$ -invariant and therefore is Γ_s -invariant. Let us consider the unitary group

$$\text{U}(V, \psi) \subset \text{GL}(V)$$

of the F -vector space V relative to ψ . A priori $\text{U}(V, \psi)$ is an algebraic group over E , but we regard it as an algebraic $\mathbf{Q}$ -group, i.e., take its Weil restriction over $\mathbf{Q}$. In particular, $\text{U}(V, \psi)(\mathbf{Q}) = \text{Aut}_F(V, \psi)$. Clearly, $\Gamma_s \subset \text{Aut}_F(V, \psi) = \text{U}(V, \psi)(\mathbf{Q})$ and therefore $G_s \subset \text{U}(V, \psi)$. The semisimplicity of G_s^0 implies that $G_s^0 \subset \text{SU}(V, \psi)$ where $\text{SU}(V, \psi)$ is the special unitary group of the F -vector space V relative to ψ . As above, we view $\text{SU}(V, \psi)$ as an algebraic $\mathbf{Q}$ -(sub)group (of $\text{U}(V, \psi)$). Clearly,

$$\text{SU}(V, \psi)(\mathbf{Q}) = \text{Aut}_F^1(V, \psi) := \{u \in \text{Aut}_F(V, \psi) \mid \det_F(u) = 1\}.$$

Since $\dim_F(V) = 2$, the centralizer $\text{End}_{\text{SU}(V, \psi)(\mathbf{Q})}(V)$ of $\text{SU}(V, \psi)(\mathbf{Q})$ in $\text{End}_{\mathbf{Q}}(V)$ (and even in $\text{End}_E(V)$) is *strictly greater* than F (see Theorem 9.2). Since the group $\text{SU}(V, \psi)(\mathbf{Q})$ is dense in $\text{SU}(V, \psi)$ with respect to Zariski topology [2, Ch. 5, Sect. 18, Cor. 18.3], the centralizer $\text{End}_{\text{SU}(V, \psi)}(V)$ of $\text{SU}(V, \psi)$ in $\text{End}_{\mathbf{Q}}(V)$ coincides with $\text{End}_{\text{SU}(V, \psi)(\mathbf{Q})}(V)$ and therefore is also *strictly greater* than F . This implies that if $G_s \subset \text{SU}(V, \psi)$ then $D_f \neq \text{End}^0(X)$. Applying Theorem 4.1, we obtain the following statement.

Lemma 10.4. *If $G_s \subset \text{SU}(V, \psi)$ then $\text{Isog}(X, \mathbf{C}(S))$ is infinite.*

End of the proof of Theorem 10.3. Since $G_s^0 \subset \text{SU}(V, \psi)$, the assertion (ii) follows readily. In order to prove the assertion (i), recall (Sect. 3.4) that $\Gamma_s^0 := \Gamma_s \cap G_s^0$ is a normal subgroup of finite index in Γ_s . Since $G_s^0 \subset \text{SU}(V, \psi)$, it follows that $\det_F(\Gamma_s) \subset \mu_F$. This implies that if we put $\Gamma^0 := \{g \in \Gamma_s \mid \det_F(g) = 1\}$ then Γ^0 is a normal subgroup in Γ_s and the quotient Γ_s/Γ^0 is a finite cyclic subgroup, whose order divides r_F . Using the construction of Section 3.8 applied to Γ^0 , we get a finite étale Galois cover $S^0 \rightarrow S$ with Galois group Γ_s/Γ^0 and abelian S^0 -scheme $\mathcal{X}^0 = \mathcal{X} \times_S S^0$ such that if $s_0 \in S^0$ lies above s then the image Γ_{s_0} of the corresponding monodromy representation coincides with Γ^0 . (Here we identify $\mathcal{X}_s$ with $\mathcal{X}_{s_0}^0$ and $H_1(\mathcal{X}_{s_0}^0, \mathbf{Q})$ with $H_1(\mathcal{X}_s, \mathbf{Q}) = V$.) Clearly, $\Gamma_{s_0} = \Gamma^0 \subset \text{SU}(V, \psi)$. Applying Lemma 10.4 to the abelian S^0 -scheme $\mathcal{X}^0 = \mathcal{X} \times_S S^0$, we conclude that $\text{Isog}(X, \mathbf{C}(S^0))$ is infinite. In order to finish the proof of (i), notice that the field extension $\mathbf{C}(S^0)/\mathbf{C}(S)$ is normal and its Galois group coincides with Γ_s/Γ^0 . $\square$

Remark 10.5. We keep the notation and assumptions of Theorem 10.3.

- (1) $\text{Hdg}(\mathcal{X}_s) = \text{U}(V, \psi)$ [18, Sect. 7.5]. It follows from Deligne's Theorem 3.5 that $G_s^0 = \text{SU}(V, \psi)$.
- (2) It follows from Theorem 9.2 that the centralizer $\text{End}_{\text{SU}(V, \psi)}(V)$ of $\text{SU}(V, \psi)$ in $\text{End}_{\mathbf{Q}}(V)$ is a four-dimensional central simple E -algebra D' containing F .
- (3) Suppose that $G_s = G_s^0$. It follows that D_f contains D' and therefore does not coincide with $F = \text{End}_S^0(\mathcal{X})$. However, the center $\mathcal{E}$ of D_f lies in F . Since $F \subset D' \subset D_f$ and the center of D' is E , we conclude that $\mathcal{E} \subset E$. This means that either $\mathcal{E} = \mathbf{Q}$ or $\mathcal{E} = E$, because E is a quadratic field. In both cases D_f is a central simple $\mathcal{E}$ -algebra. Applying Proposition 4.4.11 of [6], we conclude that $\mathcal{E} \neq \mathbf{Q}$, i.e., $\mathcal{E} = E$. Since $\dim_{\mathbf{Q}}(D_f)$ must divide 8 and $\dim_{\mathbf{Q}}(D') = 8$, we conclude that $D_f = D'$. Applying again the same Proposition, we conclude that D_f is ramified at one infinite place of E and unramified at another one. In particular, D_f is not isomorphic to the matrix algebra $M_2(E)$, i.e., D_f is a *quaternion* (division) E -algebra.

Example 10.6. Recall that the 5th cyclotomic field $F := \mathbf{Q}(\mu_5)$ is a CM-field of degree 4. Notice that μ_F is a cyclic group of order 10, i.e., $r_F = 10$. Let us consider a smooth genus zero affine curve

$$S = \mathbf{A}^1 \setminus \{0, 1\} = \mathbf{P}^1 \setminus \{0, 1, \infty\}$$

with coordinate λ and a family $\mathfrak{C} \rightarrow S$ of genus four smooth projective curves over S defined by the equation

$$y^5 = x(x-1)(x-\lambda)$$

and the corresponding family of jacobians $f : \mathcal{J} \rightarrow S$. Clearly, $\mathcal{J}$ is a (principally) polarized abelian S -scheme of relative dimension four, $\mathbf{C}(S) = \mathbf{C}(\lambda)$ is the field of rational functions in one variable over $\mathbf{C}$ and the generic fiber J of f is the jacobian of the $\mathbf{C}(\lambda)$ -curve $y^5 = x(x-1)(x-\lambda)$ of genus four. It follows from [13, Lemma 2.2 and Prop. 2.7] that $\mathcal{J}$ is *not* isotrivial and

$$\mathrm{End}^0(J) = \mathrm{End}_{\mathbf{C}(\lambda)}^0(J) = F.$$

This implies that J is absolutely simple and therefore $\mathcal{J}$ is *not* weakly isotrivial. By Theorem 10.3(i), there exists a cyclic extension $L/\mathbf{C}(\lambda)$ that has degree r dividing 10, is unramified outside $\{0, 1, \infty\}$ and such that $\mathrm{Isog}(J, L)$ is *infinite*. Using Kummer theory, we obtain easily that there are nonnegative integers a and b such that

$$0 \leq a < r \leq 10, \quad 0 \leq b < r \leq 10$$

and $L = \mathbf{C}(\lambda)(\sqrt[r]{\lambda^a(\lambda-1)^b})$. Clearly, $\mathbf{C}(\lambda) \subset L \subset \mathbf{C}(\sqrt[10]{\lambda}, \sqrt[10]{\lambda-1})$. It follows from Remark 2.3 that $\mathrm{Isog}(J, \mathbf{C}(\sqrt[10]{\lambda}, \sqrt[10]{\lambda-1}))$ is infinite. Now Corollary 4.6 implies that $\mathrm{Isog}_1(J^2, \mathbf{C}(\sqrt[10]{\lambda}, \sqrt[10]{\lambda-1}), \ell)$ is *infinite* for all but finitely many primes ℓ congruent to 1 modulo 4.

Notice that $\mathbf{C}(\sqrt[10]{\lambda}, \sqrt[10]{\lambda-1})$ is the field of the rational functions on the affine Fermat curve $S' : u^{10} - v^{10} = 1$, $u \neq 0, v \neq 0$ with $u = \sqrt[10]{\lambda}, v = \sqrt[10]{\lambda-1}$ and $S' \rightarrow S$, $\lambda = u^{10}$ is the corresponding finite étale cover.

REFERENCES

- [1] A. Borel, J-P. Serre, Théorèmes de finitude en cohomologie galoisienne, Comm. Math. Helv. 39 (1964) 111–164.
- [2] A. Borel, Linear algebraic groups (W.A. Benjamin, New York Amsterdam, 1969).
- [3] A. Borel, Some metric properties of arithmetic quotients of symmetric spaces and an extension theorem, J. Differential Geometry 6 (1972) 543–560.
- [4] S. Bosch, W. Lütkebohmert, M. Raynaud, Néron models (Springer-Verlag, Berlin Heidelberg New York, 1990).
- [5] N. Bourbaki, Algèbre, Chap. IX (Hermann, Paris, 1959).
- [6] P. Deligne, Théorie de Hodge.II, Publ. Math. IHES 40 (1971) 5–58.
- [7] P. Deligne, La conjecture de Weil pour les surfaces K3, Invent. Math. 15 (1972), 206–226.
- [8] P. Deligne, Hodge cycles on abelian varieties (notes by J. S. Milne). Lecture Notes in Math. 900, Springer-Verlag, Berlin, 1982, pp. 9–100.
- [9] G. Faltings, Arakelov’s theorem for abelian varieties, Invent. Math. 73 (1983) 337–347.
- [10] G. Faltings, Endlichkeitssätze für abelsche Varietäten über Zahlkörpern, Invent. Math. 73 (1983) 349–366; Erratum 75 (1984) 381.
- [11] G. Faltings, ‘Complements to Mordell’, Chapter VI in: G. Faltings, G. Wustholz et al., Rational points. Third edition. Aspects of Mathematics, E6. Friedr. Vieweg & Sohn, Braunschweig, 1992.
- [12] A. Grothendieck, Un théorème sur les homomorphismes de schémas abéliens, Invent. Math. 2 (1966) 59–78.
- [13] J. de Jong and R. Noot, ‘Jacobians with complex multiplications’, Arithmetic algebraic geometry (eds G. van der Geer, F. Oort and J. Steenbrink), Progress in Math., vol. 89 (Birkhäuser, 1991), pp. 177–192.
- [14] N. Katz, Nilpotent connection and monodromy theorem: Application of a result of Turrutin, Publ. Math. IHES 39 (1971) 175–232.
- [15] S. Lang, Abelian varieties. Second edition (Springer-Verlag, New York, 1983).
- [16] H. W. Lenstra, F. Oort, Yu. G. Zarhin, Abelian subvarieties, J. Algebra 180 (1996) 513–516.
- [17] D. Masser, Specializations of endomorphism rings of abelian varieties, Bull. Soc. math. France 124 (1996) 457–476.
- [18] B.J.J. Moonen, Yu.G. Zarhin, Hodge and Tate classes on simple abelian fourfolds, Duke Math. J. 77 (1995) 553–581.

- [19] B.J.J. Moonen, Yu. G. Zarhin, Hodge classes on abelian varieties of low dimension, *Math. Ann.* 315 (1999) 711–733.
- [20] L. Moret-Bailly, Pinceaux de variétés abéliennes, *Astérisque*, vol. 129 (1985).
- [21] D. Mumford, A note of Shimura’s paper “Discontinuous groups and abelian varieties”, *Math. Ann.* 181 (1969) 345–351.
- [22] D. Mumford, J. Fogarty, F. Kirwan, Geometric invariant theory, Third enlarged edition (Springer-Verlag, Berlin Heidelberg New York, 1994).
- [23] D. Mumford, Abelian varieties, Second edition, Oxford University Press, London, 1974.
- [24] A. N. Parshin, Minimal models of curves of genus 2 and homomorphisms of elliptic curves over function fields of finite characteristic, *Izv. AN SSSR ser. matem.* 36 (1972) 67–109; *Math. USSR Izv.* 6 (1972) 60–108.
- [25] R. Pierce, Associative Algebras (Springer-Verlag, Berlin Heidelberg New York, 1982).
- [26] M.M. Postnikov, Lie groups and Lie algebras (MIR, Moscow, 1986).
- [27] I. Reiner, Maximal orders (Academic Press, London, 1966).
- [28] J-P. Serre, Abelian ℓ -adic representations and elliptic curves. Second edition (Addison-Wesley, New York, 1989).
- [29] J-P. Serre, J. Tate, Good reduction of abelian varieties, *Ann. of Math.* (2) 88 (1968) 492–517.
- [30] A. Silverberg, Fields of definition for homomorphisms of abelian varieties, *J. Pure Applied Algebra* 77 (1992) 253–262.
- [31] A. Silverberg, Yu. G. Zarhin, Connectedness results for ℓ -adic representations associated to abelian varieties, *Comp. Math.* 97 (1995) 273–284.
- [32] J. Tate, Endomorphisms of Abelian varieties over finite fields, *Invent. Math.* 2 (1966) 134–144.
- [33] Yu. G. Zarhin, A finiteness theorem for isogenies of Abelian varieties over function fields of finite characteristic, *Funktsional Anal. i Prilozh.* 8 (1974), No. 4, 31–34; *Functional Anal. Appl.* 8 (1974) 301–303.
- [34] Yu. G. Zarhin, Endomorphisms of Abelian varieties over fields of finite characteristic, *Izv. Akad. Nauk SSSR ser. matem.* 39 (1975) 272–277; *Math. USSR Izv.* 9 (1975) 255–260.
- [35] Yu. G. Zarhin, Abelian varieties in characteristic P , *Mat. Zametki* 19 (1976) 393–400; *Mathematical Notes* 19 (1976) 240–244.
- [36] Yu. G. Zarhin, Endomorphisms of Abelian varieties and points of finite order in characteristic P , *Mat. Zametki* 21 (1977) 737–744; *Mathematical Notes* 21 (1978) 415–419.
- [37] Yu. G. Zarhin, Weights of simple Lie algebras in the cohomology of algebraic varieties, *Izv. Akad. Nauk SSSR Ser. Mat.* 48 (1984) 264–304; *Math. USSR Izv.* 24 (1985) 245–281.
- [38] Yu. G. Zarhin, A finiteness theorem for unpolarized Abelian varieties over number fields with prescribed places of bad reduction, *Invent. Math.* 79 (1985) 309–321.
- [39] Yu. G. Zarhin, A. N. Parshin, Finiteness problems in Diophantine geometry, *Amer. Math. Soc. Transl.* (2) 143 (1989) 35–102.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA

E-mail address: zarhin@math.psu.edu